arXiv:math/0109057v2 [math.AT] 19 Mar 2003

Multicomplexes, bounded cohomology and additivity of simplicial volume

Thilo Kuessner

Abstract

We discuss some additivity properties of the simplicial volume for manifolds with boundary: we give proofs of additivity for glueing amenable boundary components and of superadditivity for glueing amenable submanifolds of the boundary, and we discuss doubling of 3-manifolds.

The simplicial volume is a homotopy invariant of manifolds. It was defined in [3], using the l^1-norm on singular chains, as follows.
Let (X, Y) be a pair of spaces. For a relative homology class $h \in H_*(X, Y; \mathbb{R})$ define its Gromov norm $\parallel h \parallel := inf \{\sum_{i=1}^r \mid a_i \mid : \sum_{i=1}^r a_i\sigma_i \text{ represents } h\}$.
To get an invariant of a compact, connected, orientable n-manifold M, consider its fundamental class $[M, \partial M]$, that is the image of a generator of $H_n(M, \partial M; \mathbb{Z})$ under the canonical homomorphism $H_n(M, \partial M; \mathbb{Z}) \rightarrow H_n(M, \partial M; \mathbb{R})$, and define the simplicial volume of M, to be denoted $\parallel M, \partial M \parallel$, as the Gromov norm of $[M, \partial M]$. Extend this definition to non-connected manifolds by summing the simplicial volumina of the connected components.

The simplicial volume remains an invariant about which relatively little is known. This article is devoted to the study of the behaviour of simplicial volume with respect to cut and paste. That means, we are given an (n-1)-submanifold $F \subset M$ with $\partial F \subset \partial M$, and we wish to compare $\parallel M_F, \partial M_F \parallel$ to $\parallel M, \partial M \parallel$. Here, M_F denotes the manifold obtained by cutting off F, that is $M_F := \overline{M - N(F)}$ for a regular neighborhood $N(F)$ of F.

$\parallel M, \partial M \parallel$ may be strictly smaller than $\parallel M_F, \partial M_F \parallel$, as there may be fundamental cycles of M which are not the images of fundamental cycles of M_F. For example, we showed in [8] that $\parallel M_F, \partial M_F \parallel > \parallel M, \partial M \parallel$ if $int(M)$ is a hyperbolic n-manifold of finite volume, $n \geq 3$, and F is a closed geodesic hypersurface. On the other hand, somewhat counter-intuitively, $\parallel M, \partial M \parallel$ may be strictly larger than $\parallel M_F, \partial M_F \parallel$, as fundamental cycles for M_F need not fit together at the copies of F.

---

[0] 2000 Mathematics Subject Classification 57N65

For theorem 1, we consider the case that F is amenable, and prove:

Theorem 1: *Let M_1, M_2 be two compact n-manifolds, $A_1 \subset \partial M_1$ resp.$A_2 \subset \partial M_2$ (n-1)-dimensional submanifolds, $f : A_1 \to A_2$ a homeomorphism, $M = M_1 \cup_f M_2$ the glued manifold. If $\pi_1 A_1$, $\pi_1 A_2$ are amenable and f_* restricts to an isomorphism $f_* : ker\,(\pi_1 A_1 \to \pi_1 M_1) \to ker\,(\pi_1 A_2 \to \pi_1 M_2)$,*
then $\parallel M, \partial M \parallel \geq \parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel$.
If moreover A_1, A_2 are connected components of ∂M_1 resp. ∂M_2,
then $\parallel M, \partial M \parallel = \parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel$.
We prove analogous facts if A_1, A_2 are in the boundary of the same manifold M_1. Applied to 3-manifolds, theorem 1 means that simplicial volume is additive with respect to glueing along incompressible tori and superadditive with respect to glueing along incompressible annuli. In the special case that the boundary of the 3-manifolds consists of tori, Soma proved in [11] that simplicial volume is additive with respect to glueing incompressible tori or annuli. We think that our proof, apart from being a generalisation to manifolds with arbitrary boundary, should be of interest because the proof in [11] heavily relies on theorem 6.5.5. from Thurston's lecture notes, of which no published proof is available so far.

In chapter 4, we consider the special case of doubling a manifold, that is of glueing two (differently oriented) copies of M by the identity of ∂M. Here, two fundamental cycles of M, corresponding to opposite orientations, fit together at ∂M to give a fundamental cycle of DM, hence $\parallel DM \parallel \leq 2 \parallel M, \partial M \parallel$ trivially holds. We give precise conditions for 3-manifolds to satisfy the strict inequality.

Theorem 2: *Let M be a manifold of dimension $n \leq 3$. Then*
$\parallel DM \parallel < 2 \parallel M, \partial M \parallel$ if and only if $\parallel \partial M \parallel > 0$.
Theorem 2 will, using geometrization of 3-manifolds, follow from our result in [8] together with application of theorem 1 to 3-manifolds with boundary.

The article is organized as follows. Chapter 1 gives the necessary facts about multicomplexes. We usually refer to [3] where it contains complete proofs and just fix notations in a way useful for later chapters. Chapter 2 discusses treelike multicomplexes. The proved results are the same which are needed in [3] to prove results about glueing manifolds without boundary, our contribution consisting in writing complete proofs for the ideas indicated in section 3.5. of [3]. Theorem 1 is proved in chapter 3, and theorem 2 is proved in chapter 4.

It might be helpful for the reader that we give some non-rigorous motivation for the proof of theorem 1. Let us consider a toy example. We glue two manifolds M_1 and M_2 to get $M_1 \vee M_2$. (This is not a manifold but one may define a fundamental class in the obvious way.) We want to show $\parallel M_1 \vee M_2 \parallel \geq \parallel M_1 \parallel + \parallel M_2 \parallel$, thus we have to find an efficient way to map representatives of $[M_1 \vee M_2]$ to a sum of representatives of $[M_1]$ and of $[M_2]$. That means, we look for a chain map r, leftinverse to the inclusions, which maps simplices in $M_1 \vee M_2$ to simplices either in M_1 or in M_2.

The universal cover of $M_1 \vee M_2$ is a tree-like complex made from copies of $\tilde{M}_1$ and $\tilde{M}_2$. In a tree, any nondegenerate tripel of vertices has a unique central point, belonging to all three geodesics between the vertices. More generally, in a tree-like complex, one might try to construct a central simplex associated to any nondegenerate tupel of at least three points. For example, if M_1 and M_2 admitted hyperbolic metrics, one would have unique geodesics between two vertices in $\tilde{M}_1$ or $\tilde{M}_2$, hence also in the tree-like complex, and one can actually show that, for a nondegenerate tupel of vertices, the associated set of geodesics intersects the full 1-skeleton of exactly one top-dimensional simplex. This 'central' simplex belongs to a copy of $\tilde{M}_1$ or $\tilde{M}_2$. It is then easy to define r.

There is clearly no such construction for arbitrary manifolds. However, we show in section 2 that this construction can be done if M_1 and M_2 are aspherical, minimally complete multicomplexes. Such multicomplexes have in fact several features in common with hyperbolic spaces. The theorem that we actually prove in section 2 is a generalization of the above. We consider not only multicomplexes glued at one vertex, but glueing multicomplexes along an arbitrary submulticomplex with the additional condition that a suitable group G acts with certain transitivity properties on the submulticomplex along which the glueing is performed, and we do the above construction not for simplices but for G-orbits of simplices. (It might be tempting to consider the quotients with respect to the G-action to reduce the glueing to a generalized wedge. However, this would raise technical problems related to the fact that these quotients are not multicomplexes.)

It should be noted that in the case of glueing two closed manifolds, [3],3.5. avoids the use of multicomplexes by using the classifying spaces of the fundamental groups, where the corresponding constructions are easier. This construction generalizes to manifolds with boundary only if one were to consider manifolds with exactly one boundary component.

Technically, the line of argument is as follows. To any space X, one associates an aspherical, minimally complete multicomplex $K(X)$. Its simplicial bounded cohomology $H_b^*(K(X))$ is isometrically isomorphic to the singular bounded cohomology $H_b^*(X)$. Bounded cohomology is a device that admits to dualize problems about the simplicial volume. We show in 3.1., using the isometric isomorphisms between bounded cohomology groups, that the glueing problem for manifolds M_1, M_2 can be translated into an analogous problem for the aspherical, minimally complete multicomplexes $K(M_1), K(M_2)$.

If M is a manifold obtained by glueing along A, one lets act a large amenable

group $\Pi_M(A)$, which consists of all homotopy classes in M of pathes in A. This group action satisfies the transitivity properties needed to apply the results of chapter 2, i.e. to solve the glueing problem for the associated multicomplexes, and hence also for the original manifolds. In the proofs we will always have to distinguish two cases: the "amalgamated" case, where *two* manifolds are glued along parts of their boundary, and the "HNN"-case, where the glueing is performed by identifying two boundary subsets of *one* manifold.
Convention: For simplicity, we assume all manifolds to be oriented.
I would like to thank Michel Boileau and Elmar Vogt for discussions about the content of this paper.

1 Multicomplexes

1.1 Definitions

Definition 1 *: A multicomplex K consists of the following data:*
- Simplices*: a set V and, for any finite ordered subset $F=\{v_0,\ldots,v_n\}\subset V$ with $\mid F\mid\geq 2$, a (possibly empty) set I_F, such that for any permutation $\pi: F_1\rightarrow F_2$ there is a bijection $I_\pi: I_{F_1}\rightarrow I_{F_2}$,*
- Face maps*: for any finite ordered set $F=\{v_0,\ldots,v_n\}\subset V$ a family of maps $\left\{d_j: I_F\rightarrow I_{F-\{v_j\}}\right\}_{0\leq j\leq n}$.*

The elements of V are the 0-simplices or vertices of K. The pairs $\sigma=(F,i)$ with $\mid F\mid=n$ and $i\in I_F$ are the n-simplices of K. The j-th face of an n-simplex (F,i) is given by $\partial_j(F,i):=(F-\{v_j\},d_j(F))$.
Let K_j be the union of j-simplices and K^n the n-skeleton of K, that is the union $\cup_{0\leq j\leq n}K_j$.

The **geometric realization** $\mid K\mid$ of K is defined as follows:
$\mid K\mid$ is the set of pairs $\{(\lambda,i): i\in I_{F_\lambda}\}$, where $\lambda: V\rightarrow[0,1]$ are maps such that $\sum_{v\in V}\lambda(v)=1$ and $F_\lambda=\{v\in V:\lambda(v)>0\}$ is finite.
The set of all λ with a given F_λ and a given $i\in I_{F_\lambda}$ is canonically identified with a standard simplex and herits a topology via this identification. We consider the topology on $\mid K\mid$ defined such that a set is closed if its intersection with each simplex is closed.

Definition 2 *We call a multicomplex minimally complete, or m.c.m. for short, if the following holds: whenever $\sigma:\Delta\rightarrow\mid K\mid$ is a singular simplex, such that $\partial\sigma$ is a simplex of K, then σ is homotopic relative $\partial\Delta$ to a* unique *simplex in K.*

If σ is an n-simplex, its n-1-skeleton is the set $\{\partial_0\sigma, \ldots, \partial_n\sigma\}$. By recursion, we define that the n-k-skeleton σ_{n-k} of an n-simplex σ is the union of the n-k-skeleta of the simplices belonging to the n-k+1-skeleton of σ.

Definition 3 *We call a multicomplex K aspherical if all simplices $\sigma \neq \tau$ in K satisfy $\sigma_1 \neq \tau_1$.*

Orientation: Let $\pi : F_1 \rightarrow F_2$ be a permutation of finite sets and $i \in I_{F_1}$. We say that (F_1, i) and $(F_2, I_\pi(i))$ have the same orientation if π is even, and that they have different orientation if π is odd.
A submulticomplex L of a multicomplex K consists of a subset of the set of simplices closed under face maps. (K, L) is a pair of multicomplexes if K is a multicomplex and L is a submulticomplex of K. A group G acts simplicially on a pair of multicomplexes (K, L) if it acts on the set of simplices of K, mapping simplices in L to simplices in L, such that the action commutes with all face maps.

1.2 Bounded Cohomology

For a multicomplex K, let $F_j(K)$ be the R-vector space with basis the set of j-simplices of K. Let $O_j(K)$ be the subspace generated by the set $\{\sigma - sign(\pi)\,\pi(\sigma)\}$ where σ runs over all j-simplices, π runs over all permutations, and $\pi(F, i) := (\pi(F), I_\pi(i))$. Define the j-th chain group $C_j(K) = F_j(K)/O_j(K)$.
Let

$$C_a^j(K) := Hom_R(C_j(K), \mathbb{R}),$$

i.e., the group of antisymmmetric cochains.
If (K, L) is a pair of multicomplexes, we get an inclusion $C_j(L) \rightarrow C_j(K)$ and we define

$$C_a^j(K, L) := \left\{\omega \in C_a^j(K) : \omega(c) = 0 \text{ for all } c \in C_j(L)\right\}$$

with the norm

$$\parallel \omega \parallel_\infty := sup\{\omega(\sigma) : \sigma \text{ j-simplex}\}.$$

Let $C_b^j(K, L) := \{\omega \in C_a^j(K, L) : \parallel \omega \parallel_\infty < \infty\}$. The coboundary operator preserves $C_b^*(K, L)$, hence induces maps $\delta_b^i : C_b^i(K, L) \rightarrow C_b^{i+1}(X, Y)$. Define the bounded cohomology of (K, L) by $H_b^i(K, L) := ker\delta_b^i / im\delta_b^{i-1}$. $\parallel . \parallel_\infty$ induces a pseudo-norm $\parallel . \parallel$ on $H_b^*(K, L)$.
For the bounded cohomology of topological spaces, we refer to [5]. When dealing with a pair of multicomplexes (K, L), we will distinguish between $H_b^*(K, L)$ and $H_b^*(\mid K \mid, \mid L \mid)$. Since any bounded singular cochain is in particular a bounded simplicial cochain, we have an inclusion $h : C_b^*(\mid K \mid) \rightarrow C_b^*(K)$.

Proposition 1 *(Isometry lemma, [3], S.43): If K is a connected minimally complete multicomplex with infinitely many vertices, then $h^* : H_b^*(\mid K \mid) \rightarrow H_b^*(K)$ is an isometric isomorphism.*

For an n-dimensional compact, connected, orientable manifold let β_M be the unique class in $H^n(M, \partial M)$ such that $< \beta_M, [M, \partial M] >= 1$.
By [3], see [7], Thm.2.1., $\| M, \partial M \| = \frac{1}{\|\beta_M\|}$. In particular, $\| M, \partial M \| = 0$ if and only if $im\left(H_b^n(M, \partial M) \rightarrow H^n(M, \partial M)\right) = 0$.

It can be shown ([3],[5]) that the bounded cohomology and its pseudonorm depend only on the fundamental group. This works also for pairs (X, Y), if $\pi_1 Y \rightarrow \pi_1 X$ is injective. In particular, one has:

Lemma 1 *: If M and N are compact manifolds with incompressible boundary of the same dimension and there exists a map $f : (M, \partial M) \rightarrow (N, \partial N)$ inducing an isomorphism of pairs of fundamental groups, then $\| M, \partial M \| = deg(f) \| N, \partial N \|$.*

1.3 Aspherical multicomplexes

Proposition 2 *([3], p.46): Let X be a topological space. Then there is an aspherical m.c.m. $K(X)$ such that there is an isometric isomorphism $I : H_b^*(K(X)) \rightarrow H_b^*(X)$.*

Warning: It will be convenient for us to use notation different from Gromov's, since we will make use of a certain functoriality of $K(X)$. So one should be aware that our $K(X)$ corresponds to K/Γ_1 in Gromov's notation, as well as that our $\tilde{K}(X)$ will correspond to Gromov's K or $K(X)$. According to [3], $\mid K(X) \mid$ is actually an aspherical topological space. We will only need that $K(X)$ is aspherical in the sense of definition 3, what will follow from the geometric description of $K(X)$.
Because it will be of importance in the proof of lemma 8 and 9, we mention that I is constructed as the following composition:

$$H_b^*(K(X)) \xrightarrow{p^*} H_b^*\left(\tilde{K}(X)\right) \xrightarrow{(h^*)^{-1}} H_b^*\left(\mid \tilde{K}(X) \mid\right) \xrightarrow{(S^*)^{-1}} H_b^*(X).$$

Here, h^* is the isomorphism from proposition 1, p and S are described in the geometrical description below.
A proof of proposition 2 is given in [3]. Because it will be crucial for the proof of theorem 1, we recall the geometrical description of $K(X)$ as it can be read off the constructions in [3].

Geometrical description of $\tilde{K}(X)$: For a topological space X, $\tilde{K}(X)$ is the multicomplex defined as follows. Its 0-skeleton is $\{x : x \in X\}$. Its 1-skeleton is $\left\{(\{x, y\}, i) : x \neq y \in X, i \in I_{\{x,y\}}\right\}$ where $I_{\{x,y\}}$ is the set of homotopy classes relative $\{0, 1\}$ of maps from $[0, 1]$ to X mapping 0 to x and 1 to y. Having defined the n-1-skeleton of $\tilde{K}(X)$, the n-simplices of $\tilde{K}(X)$ are the pairs $(\{x_0, \ldots, x_n\}, i)$ with $x_0, \ldots, x_n \in X$ and i a homotopy class relative $\partial\Delta^n$ of mappings from the

standard simplex Δ^n to X, taking the i-th vertex of Δ^n into x_i for $i = 0, \ldots, n$ and the n-1-skeleton of Δ^n into the n-1-skeleton of $\tilde{K}(X)$.
We have a canonical continuous mapping $S : \widetilde{K}(X) \to X$ which maps each singular simplex to its image in X.

Geometrical description of $K(X)$: The multicomplex $K(X)$ is obtained from $\tilde{K}(X)$ by identifying, via simplicial maps, all n-simplices with a common n-1-skeleton, for all $n \geq 2$, succesively in order of increasing dimension.
In particular, we have a canonical projection $p : \tilde{K}(X) \to K(X)$.

Proposition 3 *: Let $Y \subset X$ be a subspace, such that $\pi_1(Y, y) \to \pi_1(X, y)$ is injective for all $y \in Y$. Then $K(Y)$ is a submulticomplex of $K(X)$.*
$I : H_b^*(K(X), K(Y)) \to H_b^*(X, Y)$ is an isometric isomorphism.

Proof: If two distinct 1-simplices in Y mapped to the same 1-simplex in X, the corresponding pathes in Y would be in $ker(\pi_1 Y \to \pi_1 X)$. By asphericity, simplices are determined by their 1-skeleton and the first claim follows. From the five lemma, I is an isomorphism. Therefore, it must be an isometry, since I and I^{-1} are composed by maps of norm ≤ 1. □

1.4 Amenable Groups and Averaging

Definition 4 *: For a group Γ let $B(\Gamma)$ be the space of bounded real-valued functions on Γ. Γ is called amenable if there is a Γ-invariant linear functional $Av : B(\Gamma) \to R$ such that $inf(f) \leq Av(f) \leq sup(f)$ holds for all $f \in B(\Gamma)$.*

If a group G acts simplicially on a pair of multicomplexes (K, L), we denote
$C_b^*\left(K^G, L^G\right) := \{c \in C_b^*(K, L) : gc = c$ for all $g \in G\}$,
$\delta_i^G := \delta \mid_{C_b^i(K^G, L^G)}$ and
$H_b^*\left(K^G, L^G\right) := ker\delta_i^G / im\delta_{i-1}^G$.

Lemma 2 *: (i) If an amenable group Γ acts on a pair of multicomplexes (K, L), and $p : C_b^*\left(K^\Gamma, L^\Gamma\right) \to C_b^*(K, L)$ is the inclusion, then there is a homomorphism $Av : H_b^*(K, L) \to H_b^*\left(K^\Gamma, L^\Gamma\right)$ such that $Av \circ p^* = id$ and $\parallel Av \parallel = 1$.*
(ii) If, moreover, all elements of Γ are homotopic to the identity in the category of continuous maps of pairs of spaces $(\mid K \mid, \mid L \mid)$, then $p^* \circ Av = id$.

Proof: : The proof of (i) works, for $L = \emptyset$, the same way as for singular bounded cohomology in [3],p.39. In the relative setting, if $L \neq \emptyset$, Av still is an isometry because of $\parallel Av \parallel \leq 1, \parallel p^* \parallel \leq 1$ and $Avp^* = id$, and it is an isomorphism as a consequence of the five lemma. Part (ii) follows from the homotopy lemma, [3],p.42 and is implicit in [3],p.46,cor.D. □

1.5 Group actions on multicomplexes

In the following, (X, Y) will be a pair of topological spaces. For a path $\gamma : [0,1] \to X$, we denote by $[\gamma]$ its homotopy class *in* X relative $\{0,1\}$.

Definition 5 *: Define* $\Pi_X(Y) :=$ $\{([\gamma_1], \ldots, [\gamma_n]) : n \in N, \gamma_1, \ldots, \gamma_n : [0,1] \to Y, \{\gamma_1(0), \ldots, \gamma_n(0)\} = \{\gamma_1(1), \ldots, \gamma_n(1)\}\}$.

$\Pi_X(Y)$ is a group with respect to the following product: $\{[\gamma_1], \ldots, [\gamma_m]\}\{[\gamma_1'], \ldots, [\gamma_n']\} := \{[\gamma_1 * \gamma_1'], \ldots, [\gamma_i * \gamma_i'], [\gamma_{i+1}], \ldots, [\gamma_m], [\gamma_1'], \ldots, [\gamma_n']\}$, where $*$ denotes the concatenation of pathes and $i \geq 0$ is chosen such that we have: $\gamma_j(1) = \gamma_j'(0)$ for $1 \leq j \leq i$ and $\gamma_j(1) \neq \gamma_k'(0)$ for $j \geq i+1, k \geq i+1$. (Such an i exists for a unique reindexing of the elements in the unordered sets $\{[\gamma_1], \ldots, [\gamma_m]\}$ and $\{[\gamma_1'], \ldots, [\gamma_n']\}$.)

Action of $\Pi_X(Y)$ on $K(X)$:
To define an action of $\Pi_X(Y)$ on the 0-skeleton of $K(X)$, we recall that $i : X \to K(X)$ maps X bijectively to $K(X)_0$. For $g = \{[\gamma_1], \ldots, [\gamma_n]\} \in \Pi_X(Y)$, we define $gi(\gamma_1(0)) = i(\gamma_1(1)), \ldots, gi(\gamma_n(0)) = i(\gamma_n(1))$ and $gi(v) = i(v)$ if $v \notin \{\gamma_1(0), \ldots, \gamma_n(0)\}$.
As a next step, we extend this to an action on the 1-skeleton of $K(X)$. Recall that 1-simplices $[\sigma]$ in $K(X)$ correspond to homotopy classes $[\sigma]$ of pathes $\sigma : [0,1] \to X$. Let $g = \{[\gamma_1], \ldots, [\gamma_n]\} \in \Pi_X(Y)$. For a 1-simplex $[\sigma]$ define $g[\sigma] = [\sigma]$ if $\sigma(0) \notin \{\gamma_1(0), \ldots, \gamma_n(0)\}$ and $\sigma(1) \notin \{\gamma_1(0), \ldots, \gamma_n(0)\}$. If $\sigma(0) = \gamma_i(0)$ and $\sigma(1)$ differs from all $\gamma_j(0)$, define $g[\sigma]$ to be the 1-simplex of $K(X)$ corresponding to the homotopy class of the concatenation $\sigma * i(\overline{\gamma_i})$, where $\overline{\gamma_i}(t) := \gamma_i(1-t)$ for $t \in [0,1]$. If $\sigma(1) = \gamma_i(0)$ and $\sigma(0)$ differs from all $\gamma_j(0)$, define $g[\sigma]$ as the 1-simplex corresponding to the homotopy class of $i(\gamma_i) * \sigma$. If $\sigma(0) = \gamma_i(0)$ and $\sigma(1) = \gamma_j(0)$, define $g\sigma$ to be the 1-simplex corresponding to the homotopy class of $i(\gamma_j) * \sigma * i(\overline{\gamma_i})$. All these definitions were independent of the choice of σ in its homotopy class relative $\{0,1\}$.
To define the action of $\Pi_X(Y)$ on all of $K(X)$, we claim that for a simplex $\sigma \in K(X)$ with 1-skeleton σ_1, and $g \in \Pi_X(Y)$, there exists *some* simplex in $K(X)$ with 1-skeleton $g\sigma_1$. Since $K(X)$ is aspherical, this will allow a unique extension of the group action from $K(X)_1$ to $K(X)$. To prove the claim, observe the following: if g is a path in X connecting v_0 to v_0' and if σ is a simplex in $K(X)$ represented by a singular simplex $\hat{\sigma}$ in X with 0-th vertex v_0', then $\hat{\sigma}$ can clearly be homotoped so that one gets a singular simplex in X with 0-th vertex v_0, leaving the other vertices fixed, so that we get a simplex whose 1-skeleton is $g\sigma_1$. Argueing succesively, we get the claim for general $g \in \Pi_X(Y)$.

Definition 6 *: Let (K, L) be a pair of multicomplexes. Let $\{e_1, \ldots, e_n\}$ and $\{e'_1, \ldots, e'_n\}$ be two n-tuples of 1-simplices in K with vertices in L. We say that $\{e_1, \ldots, e_n\}$ and $\{e'_1, \ldots, e'_n\}$ are L-related, if there are 1-simplices $f_1, \ldots, f_m$ in L such that*
- the vertices $f_1(0), f_1(1), \ldots, f_m(0), f_m(1)$ of $f_1, \ldots, f_m$ are all distinct and are in bijection with the set of vertices of $e_1, \ldots, e_n, e'_1, \ldots, e'_n$, (note that $m \leq 2n$, we do not assume the vertices of the e_i's and e'_i's to be distinct)
- the concatenations $f_k e_i f_l e'^{-1}_i$ (with f_k, f_l uniquely selected such that the vertices match) represent the identity in $\pi_1 K$.

The following observation is obvious from the construction.

Lemma 3 *: Let (X, Y) be a pair of spaces such that $\pi_1 Y \to \pi_1 X$ is injective. Then the action of $\Pi_X(Y)$ on $K(X)$ is transitive on $K(Y)$-related 1-simplices with vertices in $K(Y)$. That is, if e_1 and e_2 are $K(Y)$-related 1-simplices in $K(X)$, then exists $g \in \Pi_X(Y)$ with $ge_1 = e_2$.*

1.6 An application of averaging

Lemma 4 *: If $A \subset X$ is a subspace such that $im(\pi_1(A, x) \to \pi_1(X, x))$ is amenable for all $x \in A$, then $\Pi_X(A)$ is amenable.*

Proof: There is an exact sequence
$1 \to \bigoplus_{y \in A} im(\pi_1(A, y) \to \pi_1(X, y)) \to \Pi_X(A) \to Perm_{fin}(A) \to 1$, where $Perm_{fin}$ are the permutations with finite support.
It is well known that a group is amenable if any finitely generated subgroup is amenable. All finitely supported permutations have finite order. It follows that any finitely generated subgroup of $Perm_{fin}(A)$ is finite and therefore amenable. Also any finitely generated subgroup of $\bigoplus_{y \in Y} im(\pi_1(A, y) \to \pi_1(X, y))$ is contained in a finite sum of amenable groups and is therefore amenable. Thus $\Pi_X(A)$ is an amenable extension of an amenable group and, hence, is amenable. □

For $\epsilon \in R$ define a norm on $C_*(X, Y)$ by $\| z \|_\epsilon = \| z \| + \epsilon \| \partial z \|$. We get an induced pseudonorm $\| . \|_\epsilon$ on $H_*(X, Y)$.
More generally, if A is a union of connected components of Y, we define a norm on relative cycles of $C_*(X, Y)$ by $\| z \|_\epsilon^A = \| z \| + \epsilon \| \partial z |_A \|$ and consider the induced pseudonorm $\| . \|_\epsilon^A$ on $H_*(X, Y)$.

Proposition 4 *: If $im(\pi_1(A, y) \to \pi_1(X, y))$ is amenable for all $y \in A$, then $\| h \| = \| h \|_\epsilon^A$ for all $h \in H_*(X, Y)$.*

Proof: : With the additional assumption $A = Y$, proposition 4 becomes the equivalence theorem in [3], p.57. To get the general claim, we give a straigtforward modification of Gromov's proof.

We consider the dual norm on bounded cohomology, which, by the Hahn-Banach theorem, is induced from the dual norm on the relative cocycles. We will show that $\parallel c \parallel_\epsilon^A = \parallel c \parallel$ for relative cocycles c.

By propositions 1 and 2, we may assume that we are working with the complex of antisymmetric simplicial cochains of $K(X)$. By lemma 2, we may assume the cochains to be invariant under the action of the amenable group $\Pi_X(A)$. Hence, we may assume that the relative cocycle c factors over Q, where Q is the quotient of $F_*(K(X))/F_*(K(Y))$ under the relations $\overline{\sigma} = -\sigma$ and $a\sigma = \sigma$ for all $a \in \Pi_X(A)$ and all simplices σ, where $\overline{\sigma}$ is σ with the opposite orientation. We can define in an obvious way analogs of our norms on the dual of Q and we get then $\parallel c \parallel = \parallel c^Q \parallel$ and $\parallel c \parallel_\epsilon^A = \parallel c^Q \parallel_\epsilon^A$, where c^Q is c considered as a map from Q to R.

But in Q, any simplex σ with an edge in A becomes trivial, because there is some element of $\Pi_X(A)$ mapping σ to $\overline{\sigma}$. Hence, for any relative cycle $z \in C_*(X,Y)$, the image of $\partial z \mid_A$ in Q is trivial. Hence, $\parallel c^Q \parallel$ and $\parallel c^Q \parallel_\epsilon^A$ agree. □

Corollary 1 *: If M is a compact manifold, A a union of connected components of ∂M, and $im(\pi_1(A,x) \rightarrow \pi_1(M,x))$ is amenable for all $x \in A$, then for any $\epsilon > 0$ exists a representative z of $[M, \partial M]$ with $\parallel z \parallel \leq \parallel M, \partial M \parallel + \epsilon$ and $\parallel \partial z \mid_A \parallel \leq \epsilon$.*

2 Retraction in aspherical treelike complexes

If a group G acts simplicially on a multicomplex M, then $C_*(M)/G$ are abelian groups with well-defined boundary operator, even though M/G may not be a multicomplex. (An instructive example for the latter phenomenon is the action of $G = \Pi_X(X)$ on $K(X)$, for a topological space X.)

2.1 The 'amalgamated' case

Lemma 5 *: Assume that $(M, M'), (K, K'), (L, L')$ are pairs of path-connected, minimally complete submulticomplexes, such that*
(i) K, L are submulticomplexes of M, with inclusions $i_K : K \rightarrow M, i_L : L \rightarrow M$,
(ii) $M_0 = K_0 \cup L_0$,
(iii) $A := K \cap L$ is a submulticomplex, $\pi_1 A \rightarrow \pi_1 K$ and $\pi_1 A \rightarrow \pi_1 L$ are injective,
(iv) the inclusion $K \cup L \rightarrow M$ induces an isomorphism $\pi_1(K \cup L) \rightarrow \pi_1(M)$,
(v) K and L are aspherical in the sense of definition 3,
(vi) $K' = M' \cap K, L' = M' \cap L$.
Assume moreover that a group G acts simplicially on (M, M') such that
(vii) G maps (K, K') to (K, K') and (L, L') to (L, L'),

(viii) G acts transitively on A-related tuples of 1-simplices.
Then there is a relative chain map $r: G\backslash C_*\left(M,M'\right) \rightarrow G\backslash C_*\left(K,K'\right) \oplus G\backslash C_*\left(L,L'\right)$ in degrees $* \geq 2$ such that
- if $G\sigma$ is the orbit of a simplex in M, then $r\left(G\sigma\right)$ either is the G-orbit of a simplex in K or the G-orbit of a simplex in L,
- $ri_{K*} = id_{G\backslash C_*(K,K')}$, $ri_{L*} = id_{G\backslash C_*(L,L')}$.

Proof: We consider first the case $M' = \emptyset$.

The plan of the proof is as follows: let σ be a simplex in M, let $\tilde{\sigma}$ be a lift to the universal cover $\tilde{M}$, and let $v_0, \ldots, v_n \in \tilde{M}_0$ be the vertices of $\tilde{\sigma}$. To each pair $\left\{v_i, v_j\right\}$ we associate a family of 'minimizing' pathes $\left\{p\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right)\right\}$ parametrised by vertices $a_0^{ij}, \ldots, a_{m_{ij}}^{ij} \in A_0$ and by elements h_k^{ij} of $\pi_1 K$ or $\pi_1 L$ satisfying conditions described below. Associated to $\left\{v_0, \ldots, v_n\right\}$ and these families of 'minimizing' pathes, we construct a family of 'central' simplices $\left\{\tilde{\tau}\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right) \subset \tilde{M}\right\}$ and their projections $\left\{\tau\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right) \subset M\right\}$, which actually lie in K or L. We show then that all $\tau\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right)$, associated to a fixed $\tilde{\sigma}$, belong to the same G-orbit, and that also the simplices associated to either $g\tilde{\sigma}$ with $g \in \pi_1 M$ or to $\widetilde{g\sigma}$ with $g \in G$ belong to the same G-orbit. We define then $r\left(G\sigma\right) = G\tau$.

Assumption (iv) implies that the universal cover $\widetilde{K \cup L}$ is a submulticomplex of the universal cover $\tilde{M}$. Assumption (ii) together with assumption (iv) gives that the 0-skeleton of $\widetilde{K \cup L}$ is the whole 0-skeleton of $\tilde{M}$.
Let $\pi: \widetilde{K \cup L} \rightarrow K \cup L$ be the projection. We will need a specific section s of π on the 1-skeleton

$$s: \left(K \cup L\right)_1 \rightarrow \left(\widetilde{K \cup L}\right)_1$$
$$\sigma \rightarrow s\left(\sigma\right) = \tilde{\sigma}$$

defined as follows:

Fix a vertex $p \in A_0$ and some lift $\tilde{p} \in \tilde{A}_0 \subset \tilde{M}_0$. For any vertex v of $K \cup L$, there is some edge $e \in K_1 \cup L_1$ connecting p to v, because K and L are complete. There are unique lifts $\tilde{e}$ and $\tilde{v}$ such that $\tilde{e}$ has boundary points $\tilde{p}$ and $\tilde{v}$. This defines $\sim$ on the 0-skeleton, and also on some 1-simplices. Now, for all other 1-simplices $e \in K_1 \cup L_1$ with boundary points v and w, possibly $v = p$, we fix the unique lift $\tilde{e}$ in $\widetilde{K \cup L}$ with 0-th vertex $\tilde{v}$. (Note that in $K \cup L$ there are no edges with one vertex in $K_0 - A_0$, the other vertex in $L_0 - A_0$.)
It should be noted: if e has vertices v_0 and v_1, then $\tilde{e}$ has vertices $\tilde{v_0}$ and $h_1\tilde{v_1}$ with, a priori, $h_1 \in \pi_1\left(K \cup L\right)$. Assume that e is an edge in K. We have a unique edge connecting $\tilde{v_1}$ to $h_1\tilde{v_1}$. This edge projects to an edge f with both vertices v_1. Since v_1, as a vertex of e, belongs to K, we conclude that $f \in K_1$. This implies

$h_1 \in \pi_1 K \subset \pi_1 (K \cup L)$. In a similar way, if e is an edge in L, we conclude that $h_1 \in \pi_1 L$.
As a consequence, we get the following observation.

(A): *if* $g\tilde{e}$ *is an edge with boundary points* $h_0\tilde{v}_0$ *and* $h_1\tilde{v}_1$*, then* $g = h_0$ *and* $h_1 h_0^{-1}$ *is either* 1*, or an element of* $\pi_1 K$*, or an element of* $\pi_1 L$*.*

Indeed, we have just seen this for $g = 1$. The general case follows after applying g^{-1} to $g\tilde{e}$.
Moreover, if $g_1\tilde{e_1}$ is a 1-simplex with boundary points $h_{01}\tilde{v}_{01}$ and $h_{11}\tilde{v}_{11}$, and $g_2\tilde{e_2}$ is a 1-simplex with boundary points $h_{02}\tilde{v}_{02}$ and $h_{12}\tilde{v}_{12}$, then:

(B): *if* $g_1\tilde{e_1}$ *and* $g_2\tilde{e_2}$ *have a common boundary point* $h\tilde{v} = h_{11}\tilde{v}_{11} = h_{02}\tilde{v}_{02}$*,*
then one of the following two possibilities holds:
- $v \in A$ *or*
- $h_{01}^{-1}h_{11}$ *and* $h_{02}^{-1}h_{12}$ *either belong both to* $\pi_1 K$ *or belong both to* $\pi_1 L$*.*

Indeed, if $v \notin A$, then v is not adjacent to both, edges of K and edges of L.

Minimizing pathes:

Let v_0, v_1 be vertices of $\widetilde{K \cup L}$. By a **path** from v_0 to v_1 we mean a sequence of 1-simplices $e_1, \ldots, e_r$ such that v_0 is a vertex of e_1, e_i and e_{i+1} have a vertex in common for $1 \leq i \leq r-1$ and, v_1 is a vertex of e_r.

Given two vertices $v_1, v_2 \in \tilde{M}_0$, they belong to $\left(\widetilde{K \cup L}\right)_0$ because of (iii) and (v), and we may represent them as $v_1 = g_1\tilde{w}_1, v_2 = g_2\tilde{w}_2$ with $g_i \in \pi_1 (K \cup L)$ and $w_i \in (K \cup L)_0$ for $i = 1, 2$.

$\pi_1 (K \cup L) = \pi_1 K *_{\pi_1 A} \pi_1 L$ is an amalgamated product, hence $g_1 g_2^{-1}$ either belongs to $\pi_1 A$ or it can be decomposed as $g_1 g_2^{-1} = h_1 \ldots h_m$, where h_i are elements of $\pi_1 K - \pi_1 A$ or $\pi_1 L - \pi_1 A$ and, $h_i \in \pi_1 K$ iff $h_{i+1} \in \pi_1 L$. Such an expression is called a **normal form** of $g_1 g_2^{-1}$. If $h_1 \ldots h_m$ and $h_1' \ldots h_l'$ are two normal forms of $g_1 g_2^{-1}$, then necessarily $l = m$ and for $i = 1, \ldots, m$ belong h_i and h_i' to the same equivalence class modulo $\pi_1 A$.

We call a path $e_0, \ldots, e_{m-1}$ from $g_1\tilde{w}_1$ to $g_2\tilde{w}_2$ **minimizing** if it satisfies the following:

there is a *normal form* $g_1 g_2^{-1} = h_1 \ldots h_m$ and a *set of vertices* $a_0, \ldots, a_m \in A_0$, with $a_i \neq a_{i+1}$ for $i = 0, \ldots, m-1$, such that:
- e_0 has vertices $g_2\tilde{w}_2$ and $g_2\tilde{a}_m$
- e_i has vertices $h_{m-i+1} \ldots h_m g_2 \tilde{a}_{m-i}$ and $h_{m-i+2} \ldots h_m g_2 \tilde{a}_{m-i+1}$ for $i = 1, \ldots, m$
- e_{m+1} has vertices $g_1\tilde{w}_1$ and $g_1\tilde{a}_0$.

One should note that all these edges exist in $\widetilde{K \cup L}$ because all neighboring points project to distinct points in K resp. L and can therefore be joined by an edge in K resp. L, by completeness. Moreover, the construction should be understood such that we skip e_0 resp. e_{m+1} if $w_2 \in A$ resp. $w_1 \in A$.

It follows from (A) and (B), that these pathes are length-minimizing in the sense of being exactly the pathes with a minimum number of edges between v_1 and v_2. Since this latter characterisation depends only on v_1 and v_2, we conclude: for different sections s_1 and s_2, there is a bijection between the corresponding sets of minimizing pathes from v_1 to v_2.

Since $\tilde{K}$ and $\tilde{L}$ are universal covers of *minimally complete* multicomplexes, there is at most one edge between two vertices, and therefore a path of length m becomes uniquely determined after fixing its $m+1$ vertices. Hence, after fixing $a_0, \ldots, a_m \in A_0$ and a normal form $g_1 g_2^{-1} = h_1 \ldots h_m$, we get a unique path, to be denoted $p\left(a_0, \ldots, a_m; h_1, \ldots, h_m\right)$.

We note for later reference the following obvious observations:
(C1) *Subpathes of minimizing pathes are minimizing.*
(C2) *If* $e_1, \ldots, e_k$ *is a minimizing path,* e_k *projects to an edge in* K*,* e_{k+1} *projects to an edge in* L*,* e_k *and* e_{k+1} *have a common vertex, then* $e_1, \ldots, e_k, e_{k+1}$ *is a minimizing path.*

Intersection with simplices:

Given a set of vertices $\left\{v_i\right\}_{1\leq i\leq n}$ we consider the set

$$\bigcup_{0\leq i,j\leq n} P\left(i,j\right) = \left\{ r_{ij}^k : 1 \leq i < j \leq n, r_{ij}^k \in P\left(i,j\right) \right\},$$

where $P\left(i,j\right)$ is the set of minimizing paths from v_i to v_j.

Let $\tilde{\tau}$ be any simplex in $\widetilde{K\cup L}$. We claim that $\left\{ \tilde{\tau} \cap r_{ij}^k : r_{ij}^k \in P\left(i,j\right) \right\}$ is the full 1-skeleton of a subsimplex of $\tilde{\tau}$. We have to check the following claim:

if $\left[x,y\right], \left[z,w\right] \in \tilde{\tau}_1 \bigcap \cup_{ij} P\left(i,j\right)$*, then also* $\left[x,z\right], \left[x,w\right],$
$\left[y,z\right]$ *and* $\left[y,w\right]$ *belong to* $\tilde{\tau}_1 \bigcap \cup_{ij} P\left(i,j\right)$.

Indeed, assume that there is a minimizing path $\left\{e_0, \ldots, e_{m+1}\right\}$ with e_i having vertices $h_{m-i+1} \ldots h_m g_2 \tilde{a}_{m-i}$ and $h_{m-i+2} \ldots h_m g_2 \tilde{a}_{m-i+1}$ for $i = 1, \ldots, m$ such that $\left[x,y\right] = e_l$, that is, $x = h_{m-l+1} \ldots h_m g_2 \tilde{a}_{m-l}$ and $y = h_{m-l+2} \ldots h_m g_2 \tilde{a}_{m-l+1}$. Assume also that there is a minimizing path $\left\{e_0', \ldots, e_{m'}'\right\}$ with analogous notations such that $\left[z,w\right] = e_l'$, that is $z = h_{m'-l'+1}' \ldots h_{m'}' g_2' \tilde{a}_{m'-l'}'$ and $w = h_{m'-l'+2}' \ldots h_{m'}' g_2' \tilde{a}_{m'-l'+1}'$.

Note that all simplices in $\widetilde{K\cup L}$ project to simplices in K or in L. Assume that $\tilde{\tau}$ projects to K. By the discussion preceding observation (A), this means that h_{m-l+1} and $h_{m'-l'+1}'$ belong both to $\pi_1 K - \pi_1 A$. It follows that $h_{m-l}, h_{m-l+2}, h_{m'-l'}', h_{m'-l'+2}'$ belong to $\pi_1 L - \pi_1 A$. But this implies, for example, that $\left[x,z\right]$ is part of some minimizing path, namely the path

$$\left\{e_0, \ldots, e_{l-1}, \left[x,z\right], e_{l'-1}', \ldots, e_0'\right\}.$$

In a similar way, we see that [x,w], [y,z] and [y,w] belong to the intersection of $\tilde{\tau}$ with minimizing pathes between suitable pairs of v_i's.

The same argument works if τ projects to L.

'Central' simplices:

We are given vertices $v_0 = g_0\tilde{w}_0, \ldots, v_n = g_n\tilde{w}_n \in \tilde{M}_0$, $n \geq 2$.

We claim: if we fix, for each index pair $\{i,j\}$, a normal form $g_i g_j^{-1} = h_1^{ij} \ldots h_{m_{ij}}^{ij}$,
vertices $a_0^{ij}, \ldots, a_{m_{ij}}^{ij} \in A_0$, and the minimizing path $p_{ij} \in P(i,j)$,
then there is *at most one* **n-dimensional** simplex $\tilde{\tau} \in \widetilde{K \cup L}$
such that the intersection of $\tilde{\tau}$ with $\cup_{0 \leq i,j \leq n} p_{ij}$ is the 1-skeleton
of an **n-dimensional** simplex, i.e., is the full 1-skeleton of $\tilde{\tau}$.

(In fact, for such an n-simplex to exist, the h_{ij}'s as well as the a_{ij}'s have to satisfy obvious compatibilty conditions. We will not make use of these explicit conditions in our proof.)

We prove the claim. Assume there are two such simplices $\tilde{\tau}_1 \neq \tilde{\tau}_2$ with $dim(\tilde{\tau}_1) = dim(\tilde{\tau}_2) = n$. By assumption (vi), it suffices to show that $\tilde{\tau}_1$ and $\tilde{\tau}_2$ have the same 1-skeleta. We have to distinguish the cases that $\tilde{\tau}_1$ and $\tilde{\tau}_2$ have no common vertex or that they have a common subsimplex.

Assume first that $\tilde{\tau}_1$ and $\tilde{\tau}_2$ have no vertex in common. In the following, 'minimizing path' will mean the unique minimizing path with respect to our fixed choice of normal forms and of vertices in A. We will frequently use the following fact: each edge of $\tilde{\tau}_1$ (resp. $\tilde{\tau}_2$) is contained in the minimizing path from v_i to v_j for a unique pair $\{i,j\}$ of indices. This is true by a counting argument: there are $\frac{n(n+1)}{2}$ minimizing pathes and $\frac{n(n+1)}{2}$ edges of $\tilde{\tau}_1$, each edge belongs to some minimizing path by assumption, and no minimizing path can have two consecutive edges projecting both to K or both to L, by the definition of normal forms.
For each k and l, the minimizing path from v_k to v_l passes through $\tilde{\tau}_1$ as well as through $\tilde{\tau}_2$. Let $[w_k^1, w_l^1]$ resp. $[w_k^2, w_l^2]$ be the intersections of this minimizing path with $\tilde{\tau}_1$ resp. $\tilde{\tau}_2$.
We claim: all minimizing pathes from v_k to some v_i with $i \neq k$ pass through w_k^1 *and* w_k^2. To prove the claim, note that, by (C1), the subpath from v_k to w_k^1 is minimizing, that is, the corresponding sequence $h_1, \ldots, h_m$ is a normal form (for $\Pi_{i=1}^m h_i$) with $h_m \in \pi_1 L$ if τ_1 projects to K or vice versa. It follows from the definition of normal forms that also $\Pi_{i=1}^{m+1} h_i$ is a normal form if $h_{m+1} \in \pi_1 K$ is the element corresponding to an edge of K having w_k^1 as a vertex. Since normal forms are unique up to multiplication of the h_i's with elements of $\pi_1 A$, we conclude that there is no minimizing path from v_k to some vertex of τ_1 which does not pass through w_k^1. By the same argument, all minimizing pathes from v_k to some vertex of τ_2 have to pass through w_k^2. In particular, they have to contain the unique minimizing path from w_k^1 to w_k^2 as a subpath, by (C1). This, in turn, implies that all minimizing pathes from v_k to any $v_i, i \neq k$, contain the minimizing path from w_k^1 to w_k^2 and, in particular, contain the same edge of $\tilde{\tau}_1$. But, by the

above counting argument, it may not happen that several minimizing pathes pass through the same edge of $\tilde{\tau}_1$. This gives the contradiction.



It remains to discuss the case that $\tilde{\tau}_1$ and $\tilde{\tau}_2$ have a proper boundary face in common. Let w_k be a vertex of τ_1 which is not a vertex of τ_2 and w_l a vertex of both, τ_1 and τ_2. After reindexing, there are, by the above counting argument, v_k and v_l such that the minimizing path from v_k to v_l contains the edge $[w_k, w_l] \subset \tau_1$. By the same argument as above, all minimizing pathes from v_k to any v_i pass through w_k as well as through w_l, that is, they all contain the edge $[w_k, w_l]$ giving a contradiction. This finishes the proof of the claim. Since it will be used again in the proof that r is a chain map, we write down the following observation, which we have just proved.

(D): *If* $v_0, \ldots, v_n \in \widetilde{M}_0$ *are vertices of* M*, then their central simplex* $\tilde{\tau}$ *to some choice of* $\left\{h_k^{ij}\right\}, \left\{a_l^{ij}\right\}$*, if it exists, has vertices* $w_0, \ldots, w_n$ *such that for any* i *and* j *the minimizing path from* v_i *to* v_j *passes through* w_i *and* w_j*.*

We denote by $\tau\left(\left\{a_l^{ij}\right\}; \left\{h_k^{ij}\right\}\right)$ the projection of $\tilde{\tau}$ to $K \cup L$.

Next we claim: if we still are given $v_0 = g_0 \tilde{w_0}, \ldots, v_n = g_n \tilde{w_n} \in \tilde{M}_0$, but vary $a_0, \ldots, a_m \in A_0$ and the normal forms $g_i g_j^{-1} = h_1^{ij} \ldots h_{m_{ij}}^{ij}$, then all $\tau\left(\left\{a_l^{ij}\right\}; \left\{h_k^{ij}\right\}\right)$ belong to the same G-orbit.

Let us consider $\tau\left(\left\{a_l\right\}; \left\{h_k^{ij}\right\}\right)$ and $\tau\left(\left\{a_l'\right\}; \left\{h_k^{ij}\right\}\right)$. The same argument which showed uniqueness of $\tilde{\tau}\left(\left\{a_l\right\}; \left\{h_k^{ij}\right\}\right)$ lets us conclude that, representing the vertices of $\tilde{\tau}\left(\left\{a_l\right\}; \left\{h_k^{ij}\right\}\right)$ as $\gamma_0 \tilde{a}_0, \ldots, \gamma_n \tilde{a}_n$ and the vertices of $\tilde{\tau}\left(\left\{a_l'\right\}; \left\{h_k^{ij}\right\}\right)$ as $\gamma_0' \tilde{a_0}', \ldots, \gamma_n' \tilde{a_n}'$, we must have $\gamma_0 = \gamma_0', \ldots, \gamma_n = \gamma_n'$. Hence, corresponding edges of τ are A-related in the sense of definition 4 and belong, by assumption (viii), to the same G-orbit.

Now we consider $\tau\left(\left\{a_l\right\}; \left\{h_k^{ij}\right\}\right)$ and $\tau\left(\left\{a_l\right\}; \left\{h_k^{ij\prime}\right\}\right)$, where $g_i g_j^{-1} = h_1^{ij} \ldots h_m^{ij} = h_1^{ij\prime} \ldots h_m^{ij\prime}$ are different normal forms. Since we may argue succesively, it suffices to consider the case that different normal forms occur for only one index pair i, j. For the same reason, it suffices to consider the case that there is $1 \leq s \leq m-1$ such that $h_s^{ij\prime} = h_s^{ij} a^{-1}, h_{s+1}^{ij\prime} = a h_{s+1}^{ij}$ and $h_l^{ij\prime} = h_l^{ij}$ otherwise. Now, from the above construction, it follows that one of the following two possibilities takes place:

- if $h_l^{ij} \ldots h_m^{ij} g_2^{ij} \tilde{a}_{l-1}^{ij}$ and $h_{l+1}^{ij} \ldots h_m^{ij} g_2^{ij} \tilde{a}_l^{ij}$ are not vertices of $\tilde{\tau}\left(\left\{a_l^{ij\prime}\right\}; \left\{h_k^{ij}\right\}\right)$, then $\tilde{\tau}\left(\left\{a_l^{ij\prime}\right\}; \left\{h_k^{ij\prime}\right\}\right) = \tilde{\tau}\left(\left\{a_l^{ij\prime}\right\}; \left\{h_k^{ij}\right\}\right)$.

- if one resp. both of $h_l^{ij} \ldots h_m^{ij} g_2^{ij} \tilde{a}_{l-1}^{ij}$ and $h_{l+1}^{ij} \ldots h_m^{ij} g_2^{ij} \tilde{a}_l^{ij}$ are vertices of $\tilde{\tau}\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right)$, then $\tilde{\tau}\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij\prime}\right\}\right)$ has n resp. $n-1$ vertices in common with $\tilde{\tau}\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right)$, and has moreover as remaining vertices one or both of $h_l^{ij} \ldots h_m^{ij} g_2^{ij} \tilde{a}_{l-1}^{ij}$ and $h_{l+1}^{ij} \ldots h_m^{ij} g_2^{ij} \tilde{a}_l^{ij}$.

It follows that the 1-skeleta of $\tilde{\tau}\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right)$ and $\tilde{\tau}\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij\prime}\right\}\right)$, considered as tuples of 1-simplices, are A-related in the sense of definition 4. Indeed, that the concatenations are trivial in $\pi_1(K \cup L)$ follows from the assumption that K and L are aspherical in the sense of definition 3, which forces the concatenations to bound a 2-simplex resp. a union of two 2-simplices. By assumption (viii), we get that $\tau\left(\{a_l\};\left\{h_k^{ij}\right\}\right)$ and $\tau\left(\{a_l\};\left\{h_k^{ij\prime}\right\}\right)$ belong to the same G-orbit.

Retraction in treelike complex:

We are going to define r. Given an n-simplex $\sigma \in M$, $n \geq 2$, we consider a lift $\tilde{\sigma} \in \tilde{M}$ which projects to σ. Its vertices $v_0, \ldots v_n$ belong to $\widetilde{M}_0 = \left(\widetilde{K \cup L}\right)_0$. Now we run the above constructions with $v_0, \ldots, v_n$ and look for simplices τ with $dim(\tau) = n$.

If there is no simplex τ with $dim(\tau) = n$, we define $r(\sigma) = 0$. Otherwise, there is a unique G-orbit of simplices $\{g\tau : g \in G\}$ with $dim(g\tau) = n$ for all $g \in G$, and we define $r(G\sigma) := G\tau$.

We have to check that the definition of r does not depend on the choice of $\tilde{\sigma}$ neither on the choice of σ in its G-orbit.

Observe the following: if $\tilde{f} : \tilde{M} \to \tilde{M}$ is a simplicial self-map of the universal cover, such that $\pi\tilde{f}$ maps K to K and L to L, then $\tilde{f}$ maps minimizing pathes from v_i to v_j to minimizing pathes from $\tilde{f}(v_i)$ to $\tilde{f}(v_j)$. Hence, simplices $\tilde{\tau}$ intersecting the family of minimizing pathes associated to $v_0, \ldots, v_n$ in the full 1-skeleton of an n-simplex are mapped by $\tilde{f}$ to simplices $\tilde{f}(\tilde{\tau})$ intersecting the family of minimizing pathes associated to $\tilde{f}(v_0), \ldots, \tilde{f}(v_n)$ in the full 1-skeleton of an n-simplex. Thus, if $\tilde{\tau}$ belongs to the family of 'central' simplices associated to some simplex $\tilde{\sigma}$, then $\tilde{f}(\tilde{\tau})$ belongs to the family of 'central' simplices associated to $\tilde{f}(\tilde{\sigma})$.
We conclude: $r(G\sigma)$ does not depend on the choice of σ in its G-orbit, neither on the choice of $\tilde{\sigma}$, for fixed σ, in the orbit of the deck group.

Finally, the two desired conditions are clearly satisfied:
we have constructed $\tilde{\tau}$ with the help of the condition that it is a simplex in $\widetilde{K \cup L}$, hence $r(\sigma) = \tau$ is a simplex either in K or in L. If σ is a simplex in K or L, then clearly $r(\sigma) = \sigma$, hence r is leftinverse to i_{K*} and i_{L*}.

Compatibility with ∂-operator:

It remains to show that r is a chain map, i.e., that $\partial r(G\sigma) = r(G\partial\sigma)$ holds for all simplices σ in M.

First we consider the case $r(G\sigma) \neq 0$.
Let $v_0, \ldots, v_n$ be the vertices of a lift $\tilde{\sigma}$, and $\left\{h_k^{ij}\right\}$, $\left\{a_l^{ij}\right\}$ such that a central simplex $\tilde{\tau}$ exists. It is obvious, if we consider the set of vertices without v_k and the corresponding restricted sets of normal forms and vertices, that we get the k-th face of $\tilde{\tau}$ as a central simplex. That implies $r(G\partial_k\sigma) = \partial_k r(G\sigma)$ for all k, and hence $r(G\partial\sigma) = \partial r(G\sigma)$.

We consider now the case $r(G\sigma) = 0$.
If $r(G\partial_k\sigma) = 0$ for all faces $\partial_k\sigma$ of σ, we conclude $r(G\partial\sigma) = 0$.
So assume that for some face $\partial_k\sigma$ of σ we have $r(G\partial_k\sigma) = G\tau$ for some (n-1)-simplex τ. That means that, for the vertices $v_0, \ldots, v_{k-1}, v_{k+1}, \ldots, v_n$ and some choice of $\left\{h_m^{ij}\right\}, \left\{a_l^{ij}\right\}$ we have the central simplex $\tilde{\tau}$. Here, the h_m^{ij} and a_l^{ij} are only chosen for $i \neq k, j \neq k$. By observation (D), $\tilde{\tau}$ has vertices $w_0, \ldots, w_{k-1}, w_k, \ldots, w_n$ such that the minimizing path from v_i to w_j passes through w_i for all $i \neq k \neq j$. For $i \neq k$ consider the set P'_{ik} of the minimizing pathes from v_k to w_i. There exists some vertex v, with the property that for each i exists a path in P'_{ik} containing $[v, w_i]$ as last edge. If v were not one of $w_0, \ldots, w_{k-1}, w_{k+1}, \ldots, w_n$, then this would give us a central simplex to the vertices $v_0, \ldots, v_{k-1}, v_k, v_{k+1}, \ldots, v_n$, contradicting the assumption. Hence $v = w_j$ for some j. But this implies that the central simplices to $v_0, \ldots, v_{k-1}, v_{k+1}, \ldots, v_n$ are exactly the central simplices to $v_0, \ldots, v_{j-1}, v_{j+1}, \ldots, v_n$, and the corresponding simplices have the same orientation if and only if $j - k$ is *odd*. Therefore, $r(\partial_k\sigma)$ cancels against $r(\partial_j\sigma)$. As we find such a j for any k with $r(\partial_k\sigma) \neq 0$, we get $r(\partial\sigma) = 0$.


v_0
$\tilde{\tau}(v_0, v_1, v_2)$
$= \tilde{\tau}(v_0, v_1, v_3)$
v_2
$\tilde{\tau}(v_0, v_2, v_3)$
$= \tilde{\tau}(v_1, v_2, v_3)$
v_1
v_3

Relative construction:

Finally, we consider the general case $M' \neq \emptyset$. For the universal cover $\pi : \tilde{M} \to M$ denote the subcomplex $\tilde{M}' := \pi^{-1}(M')$. Since $A \subset M'$, the definition of 'minimizing pathes' implies that, for $x, y \in \tilde{M}'_0$, all minimizing pathes from x to y are subsets of $\tilde{M}'$. As a consequence, for $v_0, \ldots, v_n \in \tilde{M}'_0$, any central simplex, if it exists, must belong to $\tilde{M}'$. Hence, $\tau \subset M' \cap K = K'$ or $\tau \subset M' \cap L = L'$. This proves that r induces a chain map $r : C_*(M, M') \to C_*(K, K') \oplus C_*(L, L')$, finishing the proof of lemma 5. $\square$

Corollary 2 *: Let* $(M, M'), (K, K'), (L, L')$ *satisfy all assumptions of lemma 5. Let* $\gamma_1 \in H_b^p(K, K'), \gamma_2 \in H_b^p(L, L')$ *be bounded cohomology classes,* $p \geq 3$*, such* *that the action of G fixes γ_1 and γ_2. Then exists a class $\gamma \in H_b^p(M, M')$ satisfying $\| \gamma \| \leq max\{\| \gamma_1 \|, \| \gamma_2 \|\}$, such that the restrictions to K and L give back γ_1 and γ_2 and that γ is fixed by G.*

Proof: Let c_1 resp. c_2 be bounded cocycles representing γ_1 resp. γ_2. To define a bounded cocycle c, it suffices to define its value on simplices. So let σ be a simplex in M. Define:

- $c(\sigma) := c_1(r(\sigma))$ if $r(G\sigma) \in G \backslash C_*(K)$,
- $c(\sigma) := c_2(r(\sigma))$ if $r(G\sigma) \in G \backslash C_*(L)$,
- $c(\sigma) := 0$ else.

If $p \geq 3$, we get an induced map in bounded cohomology. Indeed, let $c_1 - \tilde{c_1} = \delta b_1$, $c_2 - \tilde{c_2} = \delta b_2$ for (bounded) relative (p-1)-cochains b_1, b_2 and define again $b(\tau)$ as $b_1(r(\tau)), b_2(r(\tau))$ or 0 according to whether $r(\tau)$ is in $G \backslash C_*(K)$, in $G \backslash C_*(L)$ or 0. (This definition would not work for $p = 2$.) An obvious calculation yields $c - \tilde{c} = \delta b$.

All claims of the corollary are obvious except possibly that c is indeed a relative cocycle, i.e., that δc vanishes on (p+1)-simplices in M'. Assume w.l.o.g. that $r(\sigma) \in C_*(K')$. Then $c_1(\partial r(\sigma)) = 0$ because c_1 is a relative cocycle and, therefore $\delta c(\sigma) = c(\partial \sigma) = c_1(r(\partial \sigma)) = c_1(\partial r(\sigma)) = 0$. □

2.2 The 'HNN'-case

Lemma 6 *: Assume that (K, K') is a pair of path-connected, minimally complete multicomplexes, and that A_1, A_2 are disjoint minimally complete submulticomplexes of K' such that there exists a simplicial isomorphism $F : A_1 \rightarrow A_2$. Let (L, L') be the pair of multicomplexes obtained from (K, K') by identifying σ and $F(\sigma)$ for all simplices σ in A_1 and let $P : (K, K') \rightarrow (L, L')$ be the canonical projection.*

Moreover let (M, M') be a pair of multicomplexes. Assume that

(i) L is a submulticomplex of M with inclusion $i_L : L \rightarrow M$,

(ii) $M_0 = L_0$,

(iii) $\pi_1 A_1 \rightarrow \pi_1 K$ and $\pi_1 A_2 \rightarrow \pi_1 K$ are injective,

(iv) the inclusion $L \rightarrow M$ induces an isomorphism $\pi_1 L \rightarrow \pi_1 M$,

(v) K and L are aspherical in the sense of definition 3,

(vi)$L' = M' \cap L$.

Assume moreover that a group G acts simplicially on (M, M') as well as on (K, K') such that

(vii) the action of G commutes with $i_L P$, as well as with F,

(viii) G acts transitively on j-simplices of A_1, for any $j \geq 0$.

Then there is a relative chain map $r : G \backslash C_*(M, M') \rightarrow G \backslash C_*(K, K')$ in degrees $* \geq 2$ such that

- *if* $G\sigma$ *is the* G*-orbit of a simplex in* M*, then* $r\left(G\sigma\right)$ *is the* G*-orbit of a simplex in* K*,*
- $ri_{L*}P_{*}=id_{G\backslash C_{*}(K,K')}$.

Proof: The proof of lemma 6 parallels in several aspects the proof of lemma 5. We will then be somewhat briefer in the explanations.
By (iv), the universal covering $\tilde{L}$ is a submulticomplex of $\tilde{M}$. By (ii) and (iv), $\tilde{L}_0=\tilde{M}_0$.
Denote A the image of A_1 (or A_2) in $L\subset M$. Fix a vertex $p\in A$ and some lift $\tilde{p}$. A vertex $v\in L_0$ is the image of some vertex of K, to be denoted v by abuse of notation. There is some edge in K with boundary points p and v, because K is complete. Its image under the projection P is an edge in L with boundary points p and v. There are unique lifts $\tilde{e}\in\tilde{L}_1$ and $\tilde{v}\in\tilde{L}_0$ such that $\tilde{e}$ has boundary points $\tilde{p}$ and $\tilde{v}$. If v happens to be in A, we construct $\tilde{e}$ and $\tilde{v}$ by choosing an edge e which remains in A. This is possible because A is complete.
Recall that π_1L is an HNN-extension of π_1K. We consider π_1K as subgroup of π_1L and denote by t the extending element of

$$\pi_1L=<\pi_1K,t\mid t^{-1}at=F_*a\quad\forall a\in\pi_1A>.$$

If $e\in L_1$ is an edge with boundary points v and w, we fix the unique lift $\tilde{e}\in\tilde{L}_1$ with 0-th vertex $\tilde{v}$. It will be crucial that the 1-th vertex of $\tilde{e}$ is then necessarily of the form $g\tilde{w}$ for some $w\in L_0$ with either $g\in\pi_1K$ or $g=t$. This is true because: the edge f with vertices $\tilde{w}$ and $g\tilde{w}$ projects to a closed edge in L, which is either the image of a closed edge in K or the image of an edge in K with endpoints $b\in A_1,c\in A_2$ such that $F\left(b\right)=c$. In the first case, $g\in\pi_1K$, in the second case $g=t$.

We have defined a map $\sim: L^1\rightarrow\tilde{L}^1$, satisfying
(A): *if* $g\tilde{e}$ *is an edge with boundary points* $h_0\tilde{v}_0$ *and* $h_1\tilde{v}_1$*, then* $g=h_0$ *and* $h_1h_0^{-1}$ *is either 1 or is an element of* π_1K *or equals* t*.*
Moreover, if $g_1\tilde{e_1}$ is a 1-simplex with boundary points $h_{01}\tilde{v}_{01}$ and $h_{11}\tilde{v}_{11}$, and $g_2\tilde{e_2}$ is a 1-simplex with boundary points $h_{02}\tilde{v}_{02}$ and $h_{12}\tilde{v}_{12}$, then:
(B): *if* $g_1\tilde{e_1}$ *and* $g_2\tilde{e_2}$ *have a common boundary point* $h\tilde{v}=h_{11}\tilde{v}_{11}=h_{02}\tilde{v}_{02}$*, then one of the following two possibilities holds:*
- $v\in A$ *or*
- $h_{01}^{-1}h_{11}$ *and* $h_{02}^{-1}h_{12}$ *belong both to* π_1K *or equal both* t*.*

Minimizing pathes:
Given two vertices $v_0,v_1\in\tilde{M}_0=\tilde{L}_0$, we may represent them as $v_1=g_1\tilde{w}_1,v_2=g_2\tilde{w}_2$ with $g_i\in\pi_1L$ and $w_i\in M_0$ for $i=1,2$. π_1L is an HNN-extension of π_1K,

hence $g_1g_2^{-1}$ has an expression $g_1g_2^{-1} = h_1 \ldots h_m$ with $h_i \in \pi_1 K$ or $h_i \in t$ such that $h_i \in \pi_1 K$ implies $h_{i+1} \notin \pi_1 K$. (But we allow $h_i = h_{i+1} = t$.) This expression, which we will call a **normal form**, is unique up to compatible changes of the $h_i \in \pi_1 K$ in their equivalence class modulo $\pi_1 A$.

We call a path $e_1, \ldots, e_r$ **minimizing** if there are $a_0, \ldots, a_m \in A$ and a normal form $g_1g_2^{-1} = h_1 \ldots h_m$ such that
- e_0 has vertices $g_2\tilde{w}_2$ and $g_2\tilde{a}$,
- e_i has vertices $h_{m-i+1} \ldots h_m g_2 \tilde{a}$ and $h_{m-i+2} \ldots h_m g_2 \tilde{a}$ for $i = 1, \ldots, m$,
- e_{m+1} has vertices $g_1\tilde{w}_1$ and $g_1\tilde{a}$.
One should note that the above edges exist in $\widetilde{L}$, since K is complete. The construction should be understood that we skip e_0 resp. e_{m+1} if $w_2 \in A$ resp. $w_1 \in A$.
It follows from (A) and (B), that these pathes are length-minimizing in the sense of being exactly the pathes between v_1 and v_2 with a minimum number of edges. Since this latter characterisation depends only on v_1 and v_2, we conclude: for different sections, there is a bijection between the corresponding sets of minimizing pathes from v_1 to v_2.

Since there is at most one edge between two vertices, a minimizing path becomes uniquely determined after fixing its vertices. So the only freedom in the choice of the minimizing path consists
- in the choice of $a_0, \ldots, a_m$, and
- in the choice of the h_i in their equivalence class modulo $\pi_1 A$.
The unique path corresponding to such a choice will be denoted $p\left(a_0, \ldots, a_m; h_1, \ldots, h_m\right)$.

Intersection with simplices:

Let $v_0, \ldots, v_n$ be vertices of $\tilde{L}$. Defining $P\left(i,j\right)$ like in the proof of lemma 5, we want to check that, for any simplex τ in $\tilde{L}$, $\left\{\tau \cap r_{ij}^k : r_{ij}^k \in P\left(i,j\right)\right\}$ is the full 1-skeleton of some subsimplex of τ.
Assume that $[x,y]$ and $[z,w]$ are edges of τ, with $x = h_{m-l+1} \ldots h_m g_2 \tilde{a}$, $y = h_{m-l+2} \ldots h_m g_2 \tilde{a}$, $z = h'_{m'-l'+1} \ldots h'_{m'} g'_2 \tilde{a}'$ and $w = h'_{m'-l'+2} \ldots h'_{m'} g'_2 \tilde{a}'$. By the discussion preceding observation (A), we get that h_{m-l+2} and $h'_{m'-l'+2}$ either belong both to $\pi_1 K - \pi_1 A$ or are both equal to t, and that the other of these two posibilities must hold true for $h_{m-l+1}, h_{m-l+3}, h'_{m'-l'+1}, h'_{m'-l'+3}$. This implies that $[x,z]$ is part of the minimizing path $e_0, \ldots, e_l, \left[x,z\right], e'_{l'}, \ldots, e'_0$, similarly for the other edges.

'Central' simplices:

We are given vertices $v_0 = g_0\tilde{w}_0, \ldots, v_n = g_n\tilde{w}_n \in \tilde{M}_0$, $n \geq 2$.
We fix $a_0, \ldots, a_m \in A_0$ and normal forms $g_ig_j^{-1} = h_1^{ij} \ldots h_{m_{ij}}^{ij}$. Hence, we have unique minimizing pathes p_{ij} from v_i to v_j. Then there is at most one **n-**

dimensional simplex $\tilde{\tau} \in \widetilde{L}$ such that the intersection of $\tilde{\tau}$ with $\cup_{0\leq i,j\leq n} p_{ij}$ is the 1-skeleton of an **n-dimensional** simplex, i.e., is the full 1-skeleton of $\tilde{\tau}$. This is proved by literally the same argument as in the corresponding part of the proof of lemma 5, to which we refer.
We point out that the edges of $\tilde{\tau}$ are of the form $[h_0\tilde{w}_0, h_1\tilde{w}_1]$ with $h_1h_0^{-1} \in \pi_1 K$. Indeed, the definition of minimizing pathes implies that either $h_1h_0^{-1} \in \pi_1 K$ or $h_1h_0^{-1} = t$. But the latter case would contradict the assumption $n \geq 2$, because then $[h_0\tilde{w}_0, th_0\tilde{w}_0]$ would be the only edge in $\tilde{\tau}$ having $h_0\tilde{w}_0$ as a vertex.
We consider the projection τ' of $\tilde{\tau}$ to L. By construction, the edges of τ' are projections of 1-simplices of K. Assumption (v) implies then that τ' is the projection of some simplex $\tau \in K$.
The resulting simplex in K will be denoted as $\tau\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right)$. Literally the same argument as in the proof of lemma 5 shows: if we fix $v_0,\ldots,v_n$, but vary $a_0,\ldots,a_m \in A_0$ and the normal forms $g_ig_j^{-1} = h_1^{ij}\ldots h_{m_{ij}}^{ij}$,then all $\tau\left(\left\{a_l^{ij}\right\};\left\{h_k^{ij}\right\}\right)$ belong to the same G-orbit.

Retraction in treelike complex:
Given an n-simplex $\sigma \in M$, we consider a lift $\tilde{\sigma} \in \tilde{M}$ with vertices $v_0,\ldots,v_n$, and we let $r(G\sigma) = G\tau$ if the above construction gives the G-orbit of a simplex τ with $dim(\tau) = n$, otherwise we define $r(G\sigma) = 0$. The definition of r does neither depend on the choice of the lift $\tilde{\sigma}$ nor on the choice of σ in its G-orbit, and it satisfies the two desired conditions.
The same argument as in the proof of lemma 5 shows that $r(G\partial\sigma) = G\partial r(G\sigma)$. From the assumptions follows $A \subset M'$. It is then clear from the definition that minimizing pathes between points of M' remain in M'. Thus r maps $C_*(M')$ to $C_*(K \cap P^{-1}M') = C_*(K')$. □

In an analogous manner to corollary 2, we conclude

Corollary 3 *Let* $(M,M'),(K,K'),(L,L')$ *satisfy all assumptions of lemma 6. Denote* P *the composition of the projection* $K \to L$ *with the inclusion* $L \to M$*. Let* $\gamma_1 \in H_b^p(K,K')$ *be a bounded cohomology class,* $p \geq 3$*, such that the action of* G *fixes* γ_1*. Then exists a class* $\gamma \in H_b^p(M,M')$*, satisfying* $\parallel \gamma \parallel \leq \parallel \gamma_1 \parallel$*, such that* $P^*\gamma = \gamma_1$ *and that* γ *is fixed by* G*.*

3 Glueing along amenable boundaries

3.1 Dualizing the problem

Lemma 7 *: (i): Let* M_1, M_2 *be two compact n-manifolds with boundary,* A_1, A_2 *connected (n-1)-dimensional submanifolds of* ∂M_1 *resp.* ∂M_2*,* $f: A_1 \to A_2$ *a*

homeomorphism, $M = M_1 \cup_f M_2$ *the glued manifold,* $A \subset M$ *the image of the* A_i*, and* $j_1 : (M_1, \partial M_1) \rightarrow (M, \partial M \cup A), j_2 : (M_2, \partial M_2) \rightarrow (M, \partial M \cup A)$ *the inclusions.*
Assume that the following holds: For all $\gamma_1 \in H_b^n(M_1, \partial M_1), \gamma_2 \in H_b^n(M_2, \partial M_2)$ *one can find* $\gamma \in H_b^n(M, \partial M \cup A)$ *such that* $j_1^* \gamma = \gamma_1, j_2^* \gamma = \gamma_2$ *and* $\| \gamma \| \leq max \{\| \gamma_1 \|, \| \gamma_2 \|\}$*.*
Then $\| M, \partial M \| \geq \| M_1, \partial M_1 \| + \| M_2, \partial M_2 \|$*.*
(ii): Let M_1 *be a compact n-manifold with boundary,* A_1, A_2 *disjoint connected (n-1)-dimensional submanifolds of* ∂M_1*,* $f : A_1 \rightarrow A_2$ *a homeomorphism,* $M = M_1/f$ *the glued manifold,* $A \subset M$ *the image of the* A_i*, and* $P : (M_1, \partial M_1) \rightarrow (M, \partial M \cup A)$ *the canonical projection.*
Assume that the following holds: For all $\gamma_1 \in H_b^n(M_1, \partial M_1)$ *one can find* $\gamma \in H_b^n(M, \partial M \cup A)$ *such that* $P^* \gamma = \gamma_1$ *and* $\| \gamma \| \leq \| \gamma_1 \|$*.*
Then $\| M, \partial M \| \geq \| M_1, \partial M_1 \|$*.*

Proof: (i): First consider the case that M_1 and M_2 have nontrivial simplicial volume. Then, by 1.2, the relative fundamental cocycles have preimages $\beta_1 \in H_b^n(M_1, \partial M_1)$ and $\beta_2 \in H_b^n(M_2, \partial M_2)$. Consider for $i = 1, 2$

$$\gamma_i := \| M_i, \partial M_i \| \beta_i$$

By 1.2, we have $\| \gamma_i \| = 1$.

By assumption, we get $\gamma \in H_b^n(M, \partial M \cup A)$ satisfying $\| \gamma \| \leq max \{\| \gamma_1 \|, \| \gamma_2 \|\} = 1$ and $j_1^* \gamma = \gamma_1, j_2^* \gamma = \gamma_2$.

Let $i : (M, \partial M) \rightarrow (M, \partial M \cup A)$ be the inclusion. In $H_n(M, \partial M \cup A)$ we have $i_* [M, \partial M] = j_{1*} [M_1, \partial M_1] + j_{2*} [M_2, \partial M_2]$.

Hence,

$$i^* \gamma ([M, \partial M]) = \gamma_1 ([M_1, \partial M_1]) + \gamma_2 ([M_2, \partial M_2]) = \| M_1, \partial M_1 \| + \| M_2, \partial M_2 \|$$

Thus, $\beta = \frac{1}{\| M_1, \partial M_1 \| + \| M_2, \partial M_2 \|} i^* \gamma$ is the relative fundamental cocycle of $(M, \partial M)$ and, by $\| \beta \| \leq \frac{1}{\| M_1, \partial M_1 \| + \| M_2, \partial M_2 \|}$ and duality follows $\| M, \partial M \| \geq \| M_1, \partial M_1 \| + \| M_2, \partial M_2 \|$.

Now consider the case $\| M_1, \partial M_1 \| \neq 0$ and $\| M_2, \partial M_2 \| = 0$. Consider $\gamma_2 = 0$ and again $\gamma_1 = \| M_1, \partial M_1 \| \beta_1 \in H_b^n(M_1, \partial M_1)$. Then we find $\gamma \in H_b^n(M, \partial M \cup A)$ with $j_1^* \gamma = \gamma_1, j_2^* \gamma = 0$ and $\| \gamma \| \leq \| \gamma_1 \|$. The same way as in the first case we get that $\frac{1}{\| M_1, \partial M_1 \|} i^* \gamma$ represents β and thus $\| M, \partial M \| \geq \| M_1, \partial M_1 \|$. Finally the case $\| M_1, \partial M_1 \| = \| M_2, \partial M_2 \| = 0$ is trivial anyway.
(ii): We suppose $\| M_1, \partial M_1 \| \neq 0$, since otherwise the claim is trivially true. Then the relative fundamental cocycle has preimage $\beta_1 \in H_b^n(M_1, \partial M_1)$. Consider $\gamma_1 := \| M_1, \partial M_1 \| \beta_1$. By 1.2, $\| \gamma_1 \| = 1$.

We find $\gamma \in H_b^n(M, \partial M \cup A)$ satisfying $\| \gamma \| \leq \| \gamma_1 \| = 1$ and $P^* \gamma = \gamma_1$.

Let $i : (M, \partial M) \to (M, \partial M \cup A)$ be the inclusion. In $H_n(M, \partial M \cup A)$ we have $i_*[M, \partial M] = P_*[M_1, \partial M_1]$. Hence, $i^*\gamma[M, \partial M] = \gamma P_*[M_1, \partial M_1] = P^*\gamma[M_1, \partial M_1] = \gamma_1[M_1, \partial M_1] = \parallel M_1, \partial M_1 \parallel$.
Thus, $\beta = \frac{1}{\parallel M_1, \partial M_1 \parallel} i^*\gamma$ is the relative fundamental class of $(M, \partial M)$ and by $\parallel \beta \parallel \leq \frac{1}{\parallel M_1, \partial M_1 \parallel}$ and duality follows $\parallel M, \partial M \parallel \geq \parallel M_1, \partial M_1 \parallel$. $\square$

3.2 Multicomplexes associated to glueings

The 'amalgamated' case.
We are going to consider the following situation: X_1, X_2 are topological spaces, $A_1 \subset X_1, A_2 \subset X_2$ path-connected subspaces, $f : A_1 \to A_2$ is a homeomorphism such that $f_* : \pi_1 A_1 \to \pi_1 A_2$ restricts to an isomorphism from $ker(\pi_1 A_1 \to \pi_1 X_1)$ to $ker(\pi_1 A_2 \to \pi_1 X_2)$. Let $X = X_1 \cup_f X_2$.

The assumption on f_* implies that $\pi_1 X_1, \pi_1 X_2$ inject into $\pi_1 X$. By proposition 3, it follows that
- (i) $K(X_1)$ and $K(X_2)$ are submulticomplexes of $K(X)$.
Concerning the 0-skeleta, we have
- (ii) $K(X)_0 = X = X_1 \cup X_2 = K(X_1)_0 \cup K(X_2)_0$.

Let $A = X_1 \cap X_2$ be the intersection of X_1 and X_2 as subspaces of X. There is an obvious homomorphism $\pi_1 A \to \pi_1 X$. Hence, there is a map from $K(A)$ to $K(X_1) \cap K(X_2)$, the intersection of $K(X_1)$ and $K(X_2)$ in $K(X)$, which identifies pathes in A whose composition gives an element of $ker(\pi_1 A \to \pi_1 X)$. Since this projection kills exactly the kernel of $\pi_1 K(A_i) \to \pi_1 K(X_i)$ we get that
- (iii) $\pi_1(K(X_1) \cap K(X_2))$ injects into $\pi_1 K(X_1)$ resp. $\pi_1 K(X_2)$.
In particular, $\pi_1(K(X_1) \cup K(X_2))$ is the amalgamated product of $\pi_1 K(X_1) \simeq \pi_1 X_1$ and $\pi_1 K(X_2) \simeq \pi_1 X_2$, amalgamated over $\pi_1(K(X_1) \cap K(X_2)) \simeq im(\pi_1 A_i \to \pi_1 X_i)$. But this amalgamated product is isomorphic to $\pi_1 X \simeq \pi_1 K(X)$ and we conclude
- (iv) the inclusion $K(X_1) \cup K(X_2) \to K(X)$ induces an isomorphism of fundamental groups.

Moreover, it follows from proposition 2 that
- (v) $K(X_1)$ and $K(X_2)$ are aspherical,
and we clearly have
- (vi) $K_i' = K(X_i) \cap K' \subset K(X)$ for $i = 1, 2$,
where K_i' is the image of $K(A_i)$ in $K(X_i)$ and K' is the image of $K(A)$ in $K(X)$.

Finally, from injectivity of $\pi_1 X_i \to \pi_1 X$, one gets that the canonical map $\Pi_{X_i}(A_i) \to \Pi_X(A)$ is an isomorphism for $i = 1, 2$. Consider the action of $G = \Pi_X(A)$ on $K(X)$.
- (vii) G maps $K(X_i)$ to $K(X_i)$ and K_i' to K_i' for $i = 1, 2$.
As a consequence, the action of $\Pi_X(A)$ on $K(X)$ preserves $K(X_1) \cap K(X_2) \subset K(X)$. Even though $\pi_1(X_1 \cap X_2)$ may not inject into $\pi_1 X$, analogously to lemma

3, we get:
- (viii) G acts transitively on $(K(X_1) \cap K(X_2))$-related tuples of 1-simplices in $K(X)$.

The 'HNN'-case.
Let X_1 be a topological space, A_1, A_2 path-connected subspaces of X_1, $f : A_1 \rightarrow A_2$ a homeomorphism such that f_* restricts to an isomorphism from $ker(\pi_1 A_1 \rightarrow \pi_1 X_1)$ to $ker(\pi_1 A_2 \rightarrow \pi_1 X_1)$. Let $X = X_1/_\sim$, where $x_1 \sim x_2$ if $x_1 \in A_1, x_2 \in A_2$ and $x_2 = f(x_1)$.

We have canonical, not necessarily injective, maps from $K(A_1)$ and $K(A_2)$ to $K(X_1)$. For brevity, let us denote $K_i' \subset K(X_1)$ the image of the map from $K(A_i)$, for $i = 1, 2$.
f induces a simplicial isomorphism $F : K_1' \rightarrow K_2'$. Let L be the multicomplex obtained from $K(X_1)$ by identifying σ and $F(\sigma)$ for any simplex σ in K_1'.
We have then
(i) a canonical embedding $i_L : L \rightarrow K(X)$.
Indeed, the map $K(X_1) \rightarrow K(X)$, induced by the projection, factors over L.
One checks easily:
(ii) $K(X)_0 = L_0$,
(iii) $\pi_1 K_1' \rightarrow \pi_1 K(X_1)$ and $\pi_1 K_2' \rightarrow \pi_1 K(X_1)$ are injective,
(iv) the inclusion $L \rightarrow K(X)$ induces an isomorphism of π_1's,
(v) $K(X)$ and L are aspherical in the sense of definition 3.
Let A be the image of A_1 in X and K' the image of $K(A)$ in $K(X)$. Then
(vi) the projection from $K(X_1)$ to L maps K_i' to K'.

Finally, $\pi_1 X$ is an HNN-extension of $\pi_1 X_1$, i.e. $\pi_1 X_1 \rightarrow \pi_1 X$ is injective, and one gets that the canonical map $\Pi_{X_i}(A_1) \rightarrow \Pi_X(A)$ is an isomorphism for $i = 1, 2$. Consider the action of $G = \Pi_X(A)$ on $K(X)$. We use the isomorphisms $\Pi_{X_1}(A_1) \simeq \Pi_X(A) \simeq \Pi_{X_1}(A_2)$ to get identifications with G and observe that after these identifications the action of G commutes with F. Moreover,
(vii) the action of G commutes with $i_L P$, where $P : K(X_1) \rightarrow L$ is the canonical projection.
Similarly to lemma 3, we get
(viii) G acts transitively on K'-related tuples of 1-simplices in $K(X)$.

3.3 Proof of Theorem 1

In this section, we prove superadditivity for simplicial volume of manifolds with boundary with respect to glueing along amenable subsets of the boundary. A similar result for open manifolds is the Cutting-off-theorem in [3]. One should note that, at least for manifolds with boundary, the opposite inequality need not hold. As a counterexample one may glue solid tori along disks to get a handlebody.

Lemma 8 *: (i): Let M_1, M_2 be two compact, connected n-manifolds, A_1, A_2 (n-1)-dimensional submanifolds of ∂M_1 resp.∂M_2, $f : A_1 \rightarrow A_2$ a homeomorphism and $M = M_1 \cup_f M_2$ the glued manifold.*
If f_* maps ker $(\pi_1 A_1 \rightarrow \pi_1 M_1)$ isomorphically to ker $(\pi_1 A_2 \rightarrow \pi_1 M_2)$, and if im $(\pi_1 A_1 \rightarrow \pi_1 M_1)$ is amenable, then $\parallel M, \partial M \parallel \geq \parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel$.
(ii): Let M_1 be a compact, connected n-manifold, no component of which is a 1-dimensional closed intervall, A_1, A_2 disjoint (n-1)-dimensional submanifolds of ∂M_1, $f : A_1 \rightarrow A_2$ a homeomorphism and $M = M_1 / f$ the glued manifold.
If im $(\pi_1 A_1 \rightarrow \pi_1 M_1)$ is amenable, and $f_* :$ ker $(\pi_1 A_1 \rightarrow \pi_1 M_1) \rightarrow$ ker $(\pi_1 A_2 \rightarrow \pi_1 M_1)$ is an isomorphism, then $\parallel M, \partial M \parallel \geq \parallel M_1, \partial M_1 \parallel$.

Proof: (i): For manifolds of dimensions ≤ 2 one checks easily that there is no counterexample. So we are going to assume that $n \geq 3$.
We want to check the assumption of lemma 7. We can restrict to the case that A_1 and A_2 are path-connected, since we may argue succesively for their path-connected components.

First, we make the **restrictive assumption** that $\pi_1 \partial M_i$ **and** $\pi_1 A_i$ **should inject into** $\pi_1 M_i$ for $i = 1, 2$. We will show afterwards how to handle the general case. The advantage of this assumption is that, by proposition 3, we may assume $K(\partial M_i)$ to be a submulticomplex of $K(M_i)$ and $K(\partial M \cup A)$ to be a submulticomplex of $K(M)$. Denoting by j_1, j_2, k_1, k_2 the embeddings and by I, I_1, I_2 the isometric isomorphisms from Prop.2(ii), we claim that the following diagram commutes.

$$\begin{tikzcd}
\bigoplus_{i=1}^{2} H_b^n(M_i, \partial M_i) & H_b^n(M, \partial M \cup A) \arrow[l, "j_1^* \oplus j_2^*"'] \\
\bigoplus_{i=1}^{2} H_b^n(K(M_i), K(\partial M_i)) \arrow[u, "I_1 \oplus I_2"] & H_b^n(K(M), K(\partial M \cup A)) \arrow[l, "k_1^* \oplus k_2^*"'] \arrow[u, "I"']
\end{tikzcd}$$

To see that the diagram commutes, recall that I was constructed (in section 1.3) as a composition $I = (S^*)^{-1} (h^*)^{-1} p^*$. S^* was induced by the projection $S : \widetilde{K}(X) \rightarrow X$, hence $(S_i^*)^{-1} j_i^* = j^* (S^*)^{-1}$ follows from the obvious relation $S j_i = j S_i$. Also, $h_i^* j_i^* = j^* h^*$ follows from the fact that h^* is induced by an inclusion of chain complexes. Finally, p^* is induced by the projection $p : \tilde{K}(X) \rightarrow K(X)$ described in 1.3. which clearly commutes with the inclusions coming from proposition 3.

Consider the following commutative diagram:

$$\begin{tikzcd}
\bigoplus_{i=1}^{2} H_b^n\left(K\left(M_i\right), K\left(\partial M_i\right)\right) & H_b^n\left(K\left(M\right), K\left(\partial M \cup A\right)\right) \arrow[l, "k_1^* \oplus k_2^*"'] \\
\bigoplus_{i=1}^{2} H_b^n\left(K\left(M_i\right)^{\Pi_{M_i}\left(A_i\right)}, K\left(\partial M_i\right)^{\Pi_{M_i}\left(A_i\right)}\right) \arrow[u, "p_1 \oplus p_2"] & H_b^n\left(K\left(M\right)^{\Pi_M(A)}, K\left(\partial M \cup A\right)^{\Pi_M(A)}\right) \arrow[l, "k_1^* \oplus k_2^*"'] \arrow[u, "p"']
\end{tikzcd}$$

By lemma 3, $\Pi_{M_1}\left(A_1\right), \Pi_{M_2}\left(A_2\right), \Pi_M\left(A\right)$ are amenable. Hence, by lemma 2(i), p_1, p_2, p have left inverses Av_1, Av_2, Av of norm $= 1$. For i=1,2 define

$$\gamma_i' := Av_i\left(I_i\right)^{-1}\gamma_i \in H_b^n\left(K\left(M_i\right)^{\Pi_{M_i}\left(A_i\right)}, K\left(\partial M_i\right)^{\Pi_{M_i}\left(A_i\right)}\right).$$

They satisfy $\parallel \gamma_i' \parallel = \parallel \gamma_i \parallel$.
It follows from the discussion in 3.2. that we can apply lemma 5 and corollary 2. Hence, we get

$$\gamma' \in H_b^n\left(K\left(M\right)^{\Pi_M(A)}, K_A\left(\partial M\right)^{\Pi_M(A)}\right),$$

satisfying $\parallel \gamma' \parallel \leq max\left\{\parallel \gamma_1' \parallel, \parallel \gamma_2' \parallel\right\} = max\left\{\parallel \gamma_1 \parallel, \parallel \gamma_2 \parallel\right\}$ and
$k_i^*\gamma' = \gamma_i'$ for i=1,2.
Let

$$\gamma := Ip^*\gamma' \in H_b^*\left(M, \partial M\right)$$

It satisfies $\parallel \gamma \parallel = \parallel \gamma' \parallel \leq max\left\{\parallel \gamma_1 \parallel, \parallel \gamma_2 \parallel\right\} = 1$ and
$j_1^*\gamma = j_1^*Ip^*\gamma' = I_1k_1^*p^*\gamma' = I_1p_1^*k_1^*\gamma' = I_1p_1^*Av_1\left(h_1^*\right)^{-1}\gamma_1$.

It is easy to see that $\Pi_{M_i}\left(A_i\right)$ are connected and that the actions of $\Pi_{M_i}\left(A_i\right)$ on $K\left(M_i\right)$ are continuous. Hence, all elements of $\Pi_{M_i}\left(A_i\right)$ are, as mappings from $K\left(M_i\right)$ to itself, homotopic to the identity. From part (ii) of lemma 2, we get that $p_i^* \circ Av_i = id$. Hence, we obtain $j_1^*\gamma = \gamma_1$. The same way, $j_2^*\gamma = \gamma_2$.

Thus, we have checked the assumptions of lemma 7.

We are now going to consider the **general case**, i.e., we do not assume any longer injectivity of fundamental groups.

Let $Y \subset X$ with $ker\left(\pi_1 Y \rightarrow \pi_1 X\right) \neq 0$. In this case, $\tilde{K}\left(Y\right)$ doesn't embed into $\tilde{K}\left(X\right)$. However, $\tilde{K}\left(.\right)$ is clearly functorial in the sense that continuous mappings $Y \rightarrow X$ induce simplicial maps $\tilde{K}\left(Y\right) \rightarrow \tilde{K}\left(X\right)$. In particular, the embedding induces simplicial maps $\tilde{f}: \tilde{K}\left(Y\right) \rightarrow \tilde{K}\left(X\right)$ and $f: K\left(Y\right) \rightarrow K\left(X\right)$. We consider its image $f\left(K\left(Y\right)\right)$ as a submulticomplex of $K\left(X\right)$.

We want to show that there are maps

$$H: H_b^n\left(K\left(X\right), f\left(K\left(Y\right)\right)\right) \rightarrow H_b^n\left(X, Y\right)$$

$$\overline{H} : H_b^n(X, Y) \to H_b^n(K(X), f(K(Y)))$$

of norm ≤ 1. (They are not isomorphisms, even though we will have $H\overline{H} = id$.)

To this behalf, we explain more in detail the construction of $I^{-1} = Av \circ h^* \circ S^*$ in proposition 2.

S^* is induced by a weak homotopy equivalence $S : \tilde{K}_X \to X$. From the definition of S on [3], p.42, it is clear that S maps $\tilde{f}\left(\tilde{K}_Y\right)$ to Y. Hence S induces a map S^* of norm ≤ 1 from $H_b^n(X, Y)$ to $H_b^n\left(\mid \tilde{K}_X \mid, \mid \tilde{f}\left(\tilde{K}_Y\right) \mid\right)$.

The isomorphism $Av : H_b^n\left(\tilde{K}_X\right) \to H_b^n(K(X))$ is induced by averaging over the amenable group Γ_1/Γ_n, where Γ_i is the group of simplicial automorphisms which are the identity on the i-skeleton, cf. [3], p.46. We get a map $Av : H_b^n\left(\tilde{K}_X, \tilde{f}\left(\tilde{K}_Y\right)\right) \to H_b^n(K_X, f(K_Y))$ which has norm ≤ 1 by definition of the averaging.

Finally, by proposition 1 we get an isometry $h^* : H_b^n\left(\mid \tilde{K}_X \mid, \mid \tilde{f}\left(\tilde{K}_Y\right) \mid\right) \to H_b^n\left(\tilde{K}_X, \tilde{f}\left(\tilde{K}_Y\right)\right)$. Hence, we may define $\overline{H} = Av \circ h^* \circ S^*$ and $H = j^* \circ (h^*)^{-1} \circ p^*$.

We will call H_1, H_2, H and $\overline{H_1}, \overline{H_2}, \overline{H}$ the maps corresponding to $(X, Y) = (M_1, \partial M_1), (M_2, \partial M_2)$ resp. $(M, \partial M \cup A)$.

The action of $\Pi_{A_i}(M_i)$ on $K(M_i)$ preserves $f(K(\partial M_i))$. We get the commutative diagram

$$\begin{tikzcd}
\bigoplus_{i=1}^{2} H_b^n(M_i, \partial M_i) \arrow[d, "\overline{H_1} \oplus \overline{H_2}"'] & H_b^n(M, \partial M \cup A) \arrow[l, "j_1^* \oplus j_2^*"'] \\
\bigoplus_{i=1}^{2} H_b^n(K(M_i), f(K(\partial M_i))) & H_b^n(K(M), f(K(\partial M \cup A))) \arrow[l, "k_1^* \oplus k_2^*"'] \arrow[u, "H"'] \\
\bigoplus_{i=1}^{2} H_b^n\left(K(M_i)^{\Pi_{M_i}(A_i)}, K(\partial M_i)^{\Pi_{M_i}(A_i)}\right) \arrow[u, "p_1^* \oplus p_2^*"] & H_b^n\left(K(M)^{\Pi_M(A)}, K(\partial M \cup A)^{\Pi_M(A)}\right). \arrow[l, "k_1^* \oplus k_2^*"'] \arrow[u, "p^*"']
\end{tikzcd}$$

One checks easily that all arguments in the first part of the proof go through, finishing the proof of part (i).

(ii): In dimensions ≤ 2 we check that the closed interval is the only connected counterexample. Assume then $n \geq 3$. Again we may suppose A_1, A_2 connected. We will again assume that $\pi_1(\partial M_1) \to \pi_1(M_1)$ and $\pi_1(\partial M \cup A) \to \pi_1(M)$ are

injective. The generalisation to the case of compressible boundary follows then by arguments completely analogous to those in the proof of part (i).
Like in part (i), we get a commutative diagram, where P, Q, R are the obvious projections.

$$\begin{tikzcd}
H_b^n\left(M_1, \partial M_1\right) & & H_b^n\left(M, \partial M \cup A\right) \arrow[ll, "P^*"'] \\
H_b^n\left(K\left(M_1\right), K\left(\partial M_1\right)\right) \arrow[u, "I_1"] & & H_b^n\left(K\left(M\right), K\left(\partial M \cup A\right)\right) \arrow[u, "I"] \arrow[ll, "R^*"'] \\
H_b^n\left(K\left(M_1\right)^{\Pi_{M_1}\left(A_1 \cup A_2\right)}, K\left(\partial M_1\right)^{\Pi_{M_1}\left(A_1 \cup A_2\right)}\right) \arrow[u, "p_1"] & & H_b^n\left(K\left(M\right)^{\Pi_M(A)}, K\left(\partial M\right)^{\Pi_M(A)}\right) \arrow[u, "p"] \arrow[ll, "Q^*"']
\end{tikzcd}$$

Given $\gamma_1 \in H_b^*\left(M_1, \partial M_1\right)$, define

$$\gamma_1' := Av_1 I_1^{-1} \gamma_1 \in H_b^*\left(\left(K\left(M_1\right)^{\Pi_{M_1}\left(A_1 \cup A_2\right)}, K\left(\partial M_1\right)^{\Pi_{M_1}\left(A_1 \cup A_2\right)}\right).$$

We check that the assumptions of corollary 3 are satisfied and get

$$\gamma' \in H_b^n\left(K\left(M\right)^{\Pi_M(A)}, K\left(\partial M\right)^{\Pi_M(A)}\right).$$

Then define $\gamma := Ip\gamma' \in H_b^n\left(M, \partial M \cup A\right)$. Analogously to the proof of part (i) we get that $P^*\gamma = \gamma_1$ and $\parallel \gamma \parallel \leq \parallel \gamma_1 \parallel$ holds.

Thus, we can apply lemma 7 to finish the proof. □

Lemma 9 *:*
i) Let M_1, M_2 be compact manifolds, A_1 resp. A_2 connected components of ∂M_1 resp. ∂M_2 and assume that there exist connected sets $A_i' \subset M_i$ with $A_i' \supset A_i$ and $\pi_1 A_i'$ amenable. Let $f : A_1 \rightarrow A_2$ be a homeomorphism and $M = M_1 \cup M_2 / f$ the glued manifold. Then $\parallel M, \partial M \parallel \leq \parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel$.
ii) Let M' be a compact manifold, A_1, A_2 connected components of $\partial M'$ with $\pi_1 A_i$ amenable. Let $f : A_1 \rightarrow A_2$ be a homeomorphism and $M = M'/f$ the glued manifold. Then $\parallel M, \partial M \parallel \leq \parallel M', \partial M' \parallel$.

Proof: ii) is reduced to i) via the homeomorphism $M = M' \cup_{(id,0)+(f,1)} \left(A_1 \times I\right)$. (Note that $\parallel A_1 \times I, A_1 \times \{0,1\} \parallel = 0$, since $\pi_1 A_1$ is amenable.)

To prove i), we need the following reformulation of a theorem of Matsumoto-Morita. For a space X and $q \in N$ let $C_q\left(X\right)$ be the group of singular chains and $B_q(X)$ the subgroup of boundaries. By theorem 2.8. of [9] the following two statements are equivalent:

a) there exists a number $K > 0$ such that for any boundary $z \in B_q(X)$ there is a chain $c \in C_{q+1}(X)$ satisfying $\partial c = z$ and $\parallel c \parallel < K \parallel z \parallel$,

b) the homomorphism $H_b^{q+1}(X) \to H^{q+1}(X)$ is injective.

Now let $\sum_{i=1}^m a_i\sigma_i$ and $\sum_{j=1}^n b_j\tau_j$ be representatives of $[M_1, \partial M_1]$ and $[M_2, \partial M_2]$ with

$$\sum_{i=1}^m \mid a_i \mid \leq \parallel M_1, \partial M_1 \parallel + \epsilon$$

and

$$\sum_{j=1}^n \mid b_j \mid \leq \parallel M_2, \partial M_2 \parallel + \epsilon.$$

By proposition 4 we may suppose that $\partial\left(\sum_{i=1}^m a_i\sigma_i\right) \mid_{A_1} \in C_*(\partial M_1)$ and $\partial\left(\sum_{j=1}^n b_j\tau_j\right) \mid_{A_2} \in C_*(\partial M_2)$ have norm smaller than $\frac{\epsilon}{2K}$. (Note that $\pi_1 A_i \to \pi_1 M_i$ factors over $\pi_1 A_i'$, hence has amenable image.)

Let A' be the image of A_1' in M. As $\pi_1 A'$ is amenable, $H_b^{q+1}(A') = 0$ for $q \geq 0$ ([3],[5]), hence, $H_b^{q+1}(A') \to H^{q+1}(A_i')$ is clearly injective and we get a constant K with the property in a).

Therefore, we find $c \in C_*(A') \subset C_*(M)$ with $\parallel c \parallel \leq \epsilon$ and

$$\partial c = \partial\left(\sum_{i=1}^m a_i\sigma_i + \sum_{j=1}^n b_j\tau_j\right).$$

Then $z = \sum_{i=1}^m a_i\sigma_i + \sum_{j=1}^n b_j\tau_j - c \in C_*(M_1 \cup_f M_2)$ is a fundamental cycle of norm smaller than $\parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel + 3\epsilon$. $\square$

Remark: The assumption of lemma 9 is in particular satisfied if $im\left(\pi_1\partial M_1 \to \pi_1 M_1\right)$ and $im\left(\pi_1\partial M_2 \to \pi_1 M_1\right)$ are amenable and the (singular) compression disks can be chosen to be disjoint. For 3-manifolds M_i, by a theorem of Jaco, cf. [4], there is $A_i' \subset M_i$ with $\pi_1 A_i' = im\left(\pi_1 A_i \to \pi_1 M_i\right)$ if $im\left(\pi_1 A_i \to \pi_1 M_i\right)$ is finitely presented.

Theorem 1 *:*

(i):Let M_1, M_2 be two compact n-manifolds, A_1 resp.A_2 connected components of ∂M_1 resp.∂M_2, $f: A_1 \to A_2$ a homeomorphism, $M = M_1 \cup_f M_2$ the glued manifold.

If $\pi_1 A_1$ and $\pi_1 A_2$ are amenable

and $f_*: ker\left(\pi_1 A_1 \to \pi_1 M_1\right) \to ker\left(\pi_1 A_2 \to \pi_1 M_2\right)$ is an isomorphism,

then $\parallel M, \partial M \parallel = \parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel$.

(ii): Let M_1 be a compact n-manifold, no connected component of M_1 a 1-dimensional closed intervall, A_1, A_2 connected components of ∂M_1, $f : A_1 \rightarrow A_2$ a homeomorphism, $M = M_1/f$ the glued manifold.
If $\pi_1 A_1$ is amenable and $f_* : ker\,(\pi_1 A_1 \rightarrow \pi_1 M_1) \rightarrow ker\,(\pi_1 A_2 \rightarrow \pi_1 M_1)$ is an isomorphism, then $\parallel M, \partial M \parallel = \parallel M_1, \partial M_1 \parallel$.

Proof: Theorem 1 follows from lemma 8 and 9. □

Corollary 4 *: (i) Let A be a properly embedded annulus in a compact 3-manifold M. If $im\,(\pi_1 A_1 \rightarrow \pi_1 M_A) \approx im\,(\pi_1 A_2 \rightarrow \pi_1 M_A)$ for the two images A_1, A_2 of A in M_A, then $\parallel M_A, \partial M_A \parallel \leq \parallel M, \partial M \parallel$.*
(ii) Let T be an embedded torus in a compact 3-manifold M. If
$im\,(\pi_1 T_1 \rightarrow \pi_1 M_T) \approx im\,(\pi_1 T_2 \rightarrow \pi_1 M_T)$ for the two images T_1, T_2 of T in M_T, then $\parallel M_T, \partial M_T \parallel = \parallel M, \partial M \parallel$.

Remark: If ∂M consists of tori and A is an incompressible annulus, then even $\parallel M_A, \partial M_A \parallel = \parallel M \parallel$ holds by a theorem of [11].

In [11], a version of corollary 4 has been proved for the special case that ∂M consists of tori. The principal ingredient in the proof is the following statement:
Proposition 5 ([11], Lemma 1): *Let M be a compact 3-manifold whose boundary ∂M consists of tori and H be a 3-dimensional compact submanifold of $int\,(M)$. Suppose $int\,(H)$ is a hyperbolic 3-manifold and ∂H is incompressible in M. Then we have $\parallel M, \partial M \parallel \geq \parallel H, \partial H \parallel$.*
In [12], this proposition is stated for closed M as theorem 6.5.5. without writing a proof. In [11], it is then derived for M with toral boundary, using the doubling argument, see the proof of lemma 1 in [11].
Hence, our proof seems to be the first written proof of corollary 4 and proposition 5. (It is easy to see that corollary 4 implies proposition 5.)
We want to mention that, according to Agol, an alternative proof of proposition 5 (hence, corollary 4) should be possible using the methods of [1].
We refer to [11] to see that corollary 4 actually allows to compute the simplicial volumes of all Haken 3-manifolds with (possibly empty) toral boundary.

3.4 Counterexamples

The first example shows that the condition $ker\,(\pi_1 A_1 \rightarrow \pi_1 M_1) \approx ker\,(\pi_1 A_2 \rightarrow \pi_1 M_2)$ can not be weakened.

Example 1: *Dehn fillings*
Let K be a knot in S^3 such that $S^3 - K$ admits a hyperbolic metric of finite volume. Let $V \subset S^3$ be a regular neighborhood of K. ∂V is a torus, hence, $\pi_1 \partial V$ is amenable. But

$$\parallel S^3 \parallel = 0 < \parallel S^3 - V, \partial V \parallel + \parallel V, \partial V \parallel .$$

More generally, it is known([1]) that

$$\parallel M, \partial M \parallel < \parallel M - V, \partial V \cup \partial M \parallel = \parallel M - V, \partial V \cup \partial M \parallel + \parallel V, \partial V \parallel$$

holds if both $int\,(M)$ and $int\,(M-V)$ admit a hyperbolic metric of finite volume, i.e., if M is obtained by performing hyperbolic Dehn filling at $M-V$.
Lemma 8 does not apply because the meridian of ∂V maps to zero in $\pi_1 V$, but it doesn't so in $\pi_1\left(S^3 - V\right)$ resp. $\pi_1\left(M - V\right)$.

The second example shows that the assumption 'A_i connected' in lemma 12 can not be weakened to assume only $A_i \subset A_i'$. (In particular, one can not just glue along an amenable subset $A_i \subset \partial M_i$.)

Example 2: *Heegard splittings*
Any 3-manifold can be decomposed into two handlebodies H and H', to be identified along their boundaries. Let g be the genus of H and H' and consider a set of properly embedded disks $D_1, \ldots, D_g \subset H'$ such that $H' - \cup_{i=1}^{g} V_i$ is a 3-ball B, where V_i are disjoint open regular neighborhoods of D_i. Denote $A_i = V_i \cap \partial H'$.
M is then obtained as follows: $V_1, \ldots, V_g$ are glued to H along the annuli A_i, afterwards B is glued along its whole boundary.
Of course, $\parallel V_i, \partial V_i \parallel = 0$ and $\parallel B, \partial B \parallel = 0$. Thus, if lemma 9 were applicable to the annuli A_i, we would get that $\parallel M, \partial M \parallel < \parallel H_g, \partial H_g \parallel$.
But there are 3-manifolds of arbitrarily large simplicial volume which admit Heegard splittings of a given genus. To give an explicit example, let f be a pseudo-Anosov diffeomorphism on a surface of genus g, and let M_n be the mapping tori of the iterates f^n. By Thurstons hyperbolization theorem, M_1 is hyperbolic. Hence, $\parallel M_1 \parallel > 0$ and $\parallel M_n \parallel = n \parallel M_1 \parallel$ becomes arbitrarily large. On the other hand, all M_n admit a Heegardsplitting of genus 2g+1.

4 Doubling 3-manifolds reduces simplicial volume

We recollect the theory of 3-manifolds necessary for the proof of theorem 2.

It is well-known, cf., [4] that any compact 3-manifold is a connected sum of finitely many manifolds which are irreducible or $S^2 \times S^1$, where a 3-manifold is termed irreducible if each embedded 2-sphere bounds an embedded 3-ball in M, and that any 3-manifold can be cut along finitely many disks to get pieces which have incompressible (i.e., π_1-injective) boundary.

Now let M be a compact irreducible 3-manifold with incompressible boundary. We use the Jaco-Shalen-Johannson theorem to decompose M, following [10].
In section 2 of [10], there is defined a so-called W-decomposition, which is a family of disjoint properly embedded incompressible tori and annuli. Denoting M_i the connected manifolds obtained after cutting M along the tori and annuli from the

W-decomposition, proposition 3.2. of [10] asserts that the M_i are either Seifert fibered (i.e., finitely covered by S^1-bundles), I-bundles or simple, where simple in the terminology of [10] means that any properly *immersed* incompressible torus or annulus can be isotoped into ∂M_i.

By Thurston's hyperbolization theorem, the simple pieces admit a hyperbolic metric. Moreover, we have

Proposition 6 ([13], Theorem 3): *Let M be a hyperbolic manifold such that $int\,(M)$ admits a complete hyperbolic metric. There is a totally geodesic surface in $int\,(M)$ for each non-torus component of ∂M if and only if any incompressible, properly embedded annulus is boundary-parallel and ∂M is incompressible.*

(Note that the condition "∂M incompressible", missing in [13], is necessary to exclude handlebodies. It is needed in the proof to guarantee that DM is irreducible.)

Therefore the simple pieces admit an (incomplete) hyperbolic metric such that the toral boundary components correspond to cusps and the boundary components of higher genus are totally geodesic.

For an oriented manifold M, let DM denote the double of M, defined by glueing two differently oriented copies of M via the identity of ∂M. It is trivial that $\parallel DM \parallel \leq 2 \parallel M, \partial M \parallel$. Theorem 1 implies, of course: if M is a compact 3-manifold with $\parallel \partial M \parallel = 0$, then $\parallel DM \parallel = 2 \parallel M, \partial M \parallel$. We will show that this is actually an if-and-only-if condition.

Lemma 10 *: Let N be a compact manifold of dimension ≥ 3 whose interior admits a hyperbolic metric of finite volume. Let $F \subset N$ be a closed codimension-1 submanifold which is totally geodesic for the hyperbolic metric on N. Then $\parallel N, \partial N \parallel^F > \parallel N, \partial N \parallel$.*

Proof: If $\partial N = \emptyset$, this follows from [6]. If $\partial N \neq \emptyset$, it follows from the main result of our paper [8]. □

Corollary 5 *: Let $n \geq 3$ and let M be a compact n-manifold with boundary $\partial M = \partial_0 M \cup \partial_1 M$, such that $M - \partial_0 M$ admit incomplete hyperbolic metrics of finite volume with totally geodesic boundary $\partial_1 M$.*
If $\partial_1 M \neq \emptyset$, then $\parallel DM \parallel < 2 \parallel M, \partial M \parallel$.

Proof: : Let N be the manifold obtained from two copies of M by glueing via the identity only of $\partial_1 M$. N is obtained from DM by cutting along $\partial_0 M$, hence, $\parallel N, \partial N \parallel = \parallel DM \parallel$ by theorem 1. (Note that $\pi_1 \partial_0 M$ is amenable because $\partial_0 M$ is covered by an (n-1)-torus, see p.156 in [2], for example.)
N carries a hyperbolic metric of finite volume, in which $\partial_1 M$ is totally geodesic. Hence, we get $\parallel N, \partial N \parallel < \parallel N, \partial N \parallel^{\partial_1 M} = 2 \parallel M, \partial M \parallel$. □

Theorem 2 *: If M is a compact 3-manifold with $\parallel \partial M \parallel > 0$, then $\parallel DM \parallel < 2 \parallel M, \partial M \parallel$.*

Proof: First, we want to reduce the claim to compact irreducible manifolds. For this purpose, note that $\parallel M_1 \sharp M_2, \partial \left(M_1 \sharp M_2\right) \parallel = \parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel$ holds also for manifolds with boundary (of dimension ≥ 3). Indeed, defining a fundamental class of the wedge $M_1 \vee M_2$ as $\left[M_1 \vee M_2, \partial \left(M_1 \vee M_2\right)\right] := i_{1*} \left[M_1, \partial M_1\right] + i_{2*} \left[M_2, \partial M_2\right]$, for the inclusions $i_1 : M_1 \rightarrow M_1 \vee M_2$ and $i_2 : M_2 \rightarrow M_1 \vee M_2$, we can define the simplicial volume $\parallel M_1 \vee M_2, \partial \left(M_1 \vee M_2\right) \parallel$ as the infimum over the l^1-norms of relative cycles representing $\left[M_1 \vee M_2, \partial \left(M_1 \vee M_2\right)\right]$ and it is implicit in the proof of theorem 1 that with this definition $\parallel M_1 \vee M_2, \partial \left(M_1 \vee M_2\right) \parallel = \parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel$ holds. Consider the projection from $M_1 \sharp M_2$ to $M_1 \vee M_2$ which pinches the connecting sphere to a point. It induces an isomorphism of fundamental groups (this is the point, where dimension ≥ 3 is needed) and has degree 1. By the same argument as in the proof of lemma 1, we get $\parallel M_1 \sharp M_2, \partial \left(M_1 \sharp M_2\right) \parallel = \parallel M_1 \vee M_2, \partial \left(M_1 \vee M_2\right) \parallel$.

In the same way we get generally that identification of pairs of points in manifolds doesn't change the simplicial volume. In particular one has $\parallel D \left(M_1 \sharp M_2\right) \parallel = \parallel DM_1 \parallel + \parallel DM_2 \parallel$.

Any compact 3-manifold is a connected sum of 3-manifolds which are either irreducible or $S^1 \times S^2$. Since $\parallel S^1 \times S^2 \parallel = 0$, we can reduce the claim to irreducible 3-manifolds because of $\parallel D \left(M_1 \sharp M_2\right) \parallel = \parallel DM_1 \parallel + \parallel DM_2 \parallel$ and $\parallel M_1 \sharp M_2, \partial \left(M_1 \sharp M_2\right) \parallel = \parallel M_1, \partial M_1 \parallel + \parallel M_2, \partial M_2 \parallel$.

By the discussion at the beginning of this section, we can cut M along properly embedded disks and incompressible properly embedded annuli and tori such that the obtained pieces are either Seifert fibered, I-bundles or simple, where the simple pieces admit an (incomplete) hyperbolic metric such that the toral boundary components correspond to cusps and the boundary components of higher genus are totally geodesic.

We argue that these pieces M_i satisfy the claim of theorem 2. For a Seifert fibration, the boundary consists of tori, hence, there is nothing to prove. If M_i is an I-bundle, then DM is an S^1-bundle and $\parallel DM_i \parallel = 0$ holds by corollary 6.5.3 of [12]. (But, if $\parallel \partial M_i \parallel > 0$, then $\parallel M_i \parallel \geq \frac{1}{3} \parallel \partial M_i \parallel > 0$ by the argument as in the remark on handlebodies below.) Finally, if M_i is "simple" and $\parallel \partial M_i \parallel > 0$, then the totally geodesic boundary of the hyperbolic structure is non-empty (not all boundary components can correspond to cusps), and $\parallel DM_i \parallel < 2 \parallel M_i, \partial M_i \parallel$ holds by lemma 10.

If M is a compact irreducible 3-manifold with $\parallel \partial M \parallel > 0$, then clearly $\parallel \partial M_i \parallel > 0$ holds for at least one of the pieces in its Jaco-Shalen-Johannson decomposition. To finish the proof of theorem 2, we still need the following lemma 11, where M_F is defined as in the introduction.

Lemma 11 *: Let M be a compact 3-manifold and F an incompressible, properly embedded annulus, torus or disk.*
If $\| D(M_F) \| < 2 \| M_F, \partial M_F \|$, then $\| DM \| < 2 \| M, \partial M \|$.

Proof: The claim follows from corollary 3 if F is a torus.
From now on, consider $F = A$ an annulus.
The claim will follow from the somewhat paradoxical observation that
$\| M_A, \partial M_A \| \leq \| M, \partial M \|$, but $\| D(M_A) \| \geq \| DM \|$.

DM_A is obtained from DM by cutting off an incompressible torus $T = DA$ and identifying afterwards in a different way the pairs of incompressible annuli which are ”halves” of the same copy of T. In other words $(D(M_A))_{2A} = (DM)_T$.
Look at the following picture, where dimension has been lowered by one.



Hence, we get $\| D(M_A) \| \geq \| (DM)_T \| = \| DM \|$, what implies the claim of lemma 11 because of $\| M_A, \partial M_A \| \leq \| M, \partial M \|$.
If F is a disk, the argument is the same □
This finishes the proof of theorem 2. □

Handlebodies.
Note that for any n-dimensional compact manifold holds:

$\| M, \partial M \| \geq \frac{1}{n} \| \partial M \|$. Namely, the boundary operator $\partial : H_n(M, \partial M) \rightarrow H_{n-1}(\partial M)$ maps the relative fundamental class $[M, \partial M]$ to the fundamental class $[\partial M]$. It is obvious that $\| \partial \| \leq n+1$. For a representative $\sum_{i=1}^{r} a_i \sigma_i$ of $[M, \partial M]$ one even gets $\| \partial \sum_{i=1}^{r} a_i \sigma_i \| \leq n \| \sum_{i=1}^{r} a_i \sigma_i \|$, because each σ_i has to have at least one face not in ∂M, cancelling against some other face.

Let H_g denote the 3-dimensional handlebody of genus g. H_g is a (g-1)-fold covering of H_2, hence, $\| H_g, \partial H_g \| = C(g-1)$. From the above argument follows $C \geq \frac{4}{3}$. (In fact, we showed in [7], by constructing triangulations of handlebodies, that $\frac{4}{3} \leq C \leq 3$.)

The double of H_g is the g-fold connected sum $S^2 \times S^1 \sharp \ldots \sharp S^2 \times S^1$, whose simplicial volume vanishes. This shows that there may be manifolds M of arbitrarily large simplicial volume with $\| DM \| = 0$.

In fact, we can give a precise condition when $\| DM \| = 0$ holds for a compact 3-manifold M. Recall that any compact 3-manifold can be cut along disks into finitely many pieces M_j which have incompressible boundary. We claim that $\| DM \| = 0$ if and only if all these M_j have a Jaco-Shalen-Johannson-decomposition without simple pieces in the sense of [10].

Namely, if M_j has a JSJ-decomposition without simple pieces, one easily gets $\| DM_j \| = 0$, thus, DM_j is a graph manifold by [11]. DM is obtained from $\cup DM_j$ by cutting off some 3-balls and identifying their boundaries in pairs. By the same argument as in the proof of lemma 4 in [11], DM is then a graph manifold and $\| DM \| = 0$.

To prove the other implication, assume that M_j had a simple piece H, on which we put a hyperbolic metric with totally geodesic boundary. Inside DM this gives us a submanifold H', obtained from two copies of H by glueing via the identity on *a submanifold of* ∂H. Clearly, H' admits a hyperbolic metric with totally geodesic boundary. From proposition 5, one can conclude $\| DM \| > 0$.

References

[1] I. AGOL, 'Topology of hyperbolic 3-manifolds', PhD thesis, UCSD, 1998.

[2] R. BENEDETTI and C. PETRONIO, *Lectures on Hyperbolic Geometry* (Universitext, Springer-Verlag, Berlin, 1992).

[3] M. GROMOV, 'Volume and Bounded Cohomology', *Public. Math. IHES* 56 (1982) 5-100.

[4] J. HEMPEL, *3-Manifolds* (Princeton University Press, 1976).

[5] N. IVANOV, 'Foundations of the theory of bounded cohomology', *J. Sov. Math.* 37 (1987) 1090-1114.

[6] D. JUNGREIS, 'Chains that realize the Gromov invariant of hyperbolic manifolds', *Ergodic Theory and Dynamical Systems* 17 (1997) 643-648.

[7] T. KUESSNER, 'Gromov volume of compact manifolds', diploma thesis, FU Berlin, 1996; http://www.mathematik.uni-tuebingen.de∼thilo/dipn.ps

[8] T. KUESSNER, 'Efficient fundamental cycles of cusped hyperbolic manifolds', preprint; http://front.math.ucdavis.edu/math.GT/0007003

[9] S. MATSUMOTO and S. MORITA, 'Bounded Cohomology of Certain Groups of Homeomorphisms', *Proc. Amer. Math. Soc.* 94 (1985) 539-544.

[10] W. NEUMANN and G. SWARUP, 'Canonical decompositions of 3-manifolds', *Geometry and Topology* 1 (1997) 21-40.

[11] T. SOMA, 'The Gromov invariant of links', *Invent. Math.* 64 (1981) 445-454.

[12] W. THURSTON, 'The Geometry and Topology of 3-Manifolds', lecture notes; http://msri.org/publications/books/gt3m

[13] W. THURSTON, 'Hyperbolic geometry and 3-manifolds', *Low-dimensional topology*, Proc. Conf., Bangor/Engl. 1979, Vol. 1, London Math. Soc. Lecture Note Series 48 (1982), pp.9-25.